


Universitext

Bernt Øksendal
Agnès Sulem

Applied Stochastic Control of Jump Diffusions

Second Edition



Springer

Universitext

Bernt Øksendal · Agnès Sulem

Applied Stochastic Control of Jump Diffusions

2nd Edition

With 27 Figures

 Springer

Bernt Øksendal
University of Oslo
Department of Mathematics
0316 Oslo
Norway
e-mail: oksendal@math.uio.no

Angnès Sulem
INRIA Rocquencourt
Domaine de Voluceau
78153 Le Chesnay CX
France
e-mail: agnes.sulem@inria.fr

Mathematics Subject Classification (2000): 93E20, 60G40, 60G51, 49L25, 65MXX, 47J20, 49J40, 91B28

Library of Congress Control Number: 2007921874

ISBN-10: 3-540-69825-6 Springer Berlin Heidelberg New York
ISBN-13: 978-3-540-69825-8 Springer Berlin Heidelberg New York
ISBN-13: 978-3-540-14023-8 1st ed. Springer Berlin Heidelberg New York

This work is subject to copyright. All rights are reserved, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilm or in any other way, and storage in data banks. Duplication of this publication or parts thereof is permitted only under the provisions of the German Copyright Law of September 9, 1965, in its current version, and permission for use must always be obtained from Springer. Violations are liable for prosecution under the German Copyright Law.

Springer is a part of Springer Science+Business Media
springer.com
© Springer-Verlag Berlin Heidelberg 2004, 2007

The use of general descriptive names, registered names, trademarks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

Cover design: WMXDesign, Heidelberg
Typesetting by the authors and SPi using a Springer L^AT_EX macro package

Printed on acid-free paper SPIN: 11872405 41/2141/SPi 5 4 3 2 1 0

To my family

Eva, Elise, Anders, and Karina

B. Ø.

A tous ceux qui m'accompagnent

A. S.

Preface to the Second Edition

In this second edition, we have added a chapter on optimal control of random jump fields (solutions of stochastic *partial* differential equations) and partial information control (Chap. 10). We have also added a section on optimal stopping with delayed information (Sect. 2.3). It has always been our intention to give a contemporary presentation of applied stochastic control, and we hope that the addition of these recent developments will contribute in this direction.

We have also made a number of corrections and other improvements, many of them based on helpful comments from our readers. In particular, we would like to thank Andreas Kyprianou for his valuable communications. We are also grateful to (in alphabetical order) Knut Aase, Jean-Philippe Chancelier, Inga Eide, Emil Framnes, Arne-Christian Lund, Jose-Luis Menaldi, Tamás K. Papp, Atle Seierstad, and Jens Arne Sukkestad for pointing out errors and suggesting improvements. Our special thanks go to Martine Verneuille for her skillful typing.

Oslo and Paris, November 2006

Bernt Øksendal and Agnès Sulem

Preface of the First Edition

Jump diffusions are solutions of stochastic differential equations driven by Lévy processes. Since a Lévy process $\eta(t)$ can be written as a linear combination of t , a Brownian motion $B(t)$ and a pure jump process, jump diffusions represent a natural and useful generalization of Itô diffusions. They have received a lot of attention in the last years because of their many applications, particularly in economics.

There exist today several excellent monographs on Lévy processes. However, very few of them – if any – discuss the optimal control, optimal stopping, and impulse control of the corresponding jump diffusions, which is the subject of this book. Moreover, our presentation differs from these books in that it emphasizes the applied aspect of the theory. Therefore, we focus mostly on useful verification theorems and we illustrate the use of the theory by giving examples and exercises throughout the text. Detailed solutions of some of the exercises are given at the end of the book. The exercises to which a solution is provided, are marked with an asterix *. It is our hope that this book will fill a gap in the literature and that it will be a useful text for students, researchers, and practitioners in stochastic analysis and its many applications. Although most of our results are motivated by examples in economics and finance, the results are general and can be applied in a wide variety of situations. To emphasize this, we have also included examples in biology and physics/engineering.

This book is partially based on courses given at the Norwegian School of Economics and Business Administration (NHH) in Bergen, Norway, during the Spring semesters 2000 and 2002, at INSEA in Rabat, Morocco in September 2000, at Odense University in August 2001 and at ENSAE in Paris in February 2002.

Oslo and Paris, August 2004

Bernt Øksendal and Agnès Sulem

Acknowledgments We are grateful to many people who in various ways have contributed to these lecture notes. In particular, we thank Knut Aase, Fred Espen Benth, Jean-Philippe Chancelier, Rama Cont, Hans Marius Eikseth, Nils Christian Framstad, Jørgen Haug, Monique Jeanblanc, Kenneth Karlsen, Arne-Christian Lund, Thilo Meyer-Brandis, Cloud Makasu, Sure Mataramvura, Peter Tankov, and Jan Ubøe for their valuable help. We also thank Francesca Biagini for useful comments and suggestions to the text and her detailed solutions of some of the exercises. We are grateful to Dina Haraldsson and Martine Verneuille for proficient typing and Eivind Brodal for his kind assistance. We acknowledge with gratitude the support by the French–Norwegian cooperation project Stochastic Control and Applications, Aur 99-050.

Oslo and Paris, August 2004

Bernt Øksendal and Agnès Sulem

Contents

1	Stochastic Calculus with Jump Diffusions	1
1.1	Basic Definitions and Results on Lévy Processes	1
1.2	The Itô Formula and Related Results	6
1.3	Lévy Stochastic Differential Equations	10
1.4	The Girsanov Theorem and Applications	12
1.5	Application to Finance	19
1.6	Exercises	21
2	Optimal Stopping of Jump Diffusions	27
2.1	A General Formulation and a Verification Theorem	27
2.2	Applications and Examples	31
2.3	Optimal Stopping with Delayed Information	36
2.4	Exercises	42
3	Stochastic Control of Jump Diffusions	45
3.1	Dynamic Programming	45
3.2	The Maximum Principle	52
3.3	Application to Finance	57
3.4	Exercises	61
4	Combined Optimal Stopping and Stochastic Control of Jump Diffusions	65
4.1	Introduction	65
4.2	A General Mathematical Formulation	66
4.3	Applications	71
4.4	Exercises	75
5	Singular Control for Jump Diffusions	77
5.1	An Illustrating Example	77
5.2	A General Formulation	79

5.3	Application to Portfolio Optimization with Transaction Costs	85
5.4	Exercises	87
6	Impulse Control of Jump Diffusions	91
6.1	A General Formulation and a Verification Theorem	91
6.2	Examples	95
6.3	Exercises	103
7	Approximating Impulse Control by Iterated Optimal Stopping	107
7.1	Iterative Scheme	107
7.2	Examples	117
7.3	Exercises	122
8	Combined Stochastic Control and Impulse Control of Jump Diffusions	123
8.1	A Verification Theorem	123
8.2	Examples	126
8.3	Iterative Methods	131
8.4	Exercises	132
9	Viscosity Solutions	135
9.1	Viscosity Solutions of Variational Inequalities	136
9.1.1	Uniqueness	138
9.2	The Value Function is Not Always C^1	139
9.3	Viscosity Solutions of HJBQVI	142
9.4	Numerical Analysis of HJBQVI	153
9.4.1	Finite Difference Approximation	153
9.4.2	A Policy Iteration Algorithm for HJBQVI	156
9.5	Exercises	159
10	Optimal Control of Random Jump Fields and Partial Information Control	161
10.1	A Motivating Example	161
10.2	The Maximum Principle	162
10.3	The Arrow Condition	168
10.3.1	Return to Example 10.1	169
10.4	Controls Which do not Depend on x	174
10.5	Connection with Partial Observation Control	176
10.6	Exercises	180
11	Solutions of Selected Exercises	183
11.1	Exercises of Chapter 1	183
11.2	Exercises of Chapter 2	189
11.3	Exercises of Chapter 3	202

11.4 Exercises of Chapter 4.....	209
11.5 Exercises of Chapter 5.....	211
11.6 Exercises of Chapter 6.....	215
11.7 Exercises of Chapter 7.....	227
11.8 Exercises of Chapter 8.....	232
11.9 Exercises of Chapter 9.....	235
11.10 Exercises of Chapter 10.....	240
References	243
Notation and Symbols	251
Index	255

Stochastic Calculus with Jump Diffusions

1.1 Basic Definitions and Results on Lévy Processes

In this chapter we present the basic concepts and results needed for the applied calculus of jump diffusions. Since there are several excellent books which give a detailed account of this basic theory, we will just briefly review it here and refer the reader to these books for more information.

Definition 1.1. Let $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \geq 0}, P)$ be a filtered probability space. An \mathcal{F}_t -adapted process $\{\eta(t)\}_{t \geq 0} = \{\eta_t\}_{t \geq 0} \subset \mathbb{R}$ with $\eta_0 = 0$ a.s. is called a Lévy process if η_t is continuous in probability and has stationary and independent increments.

Theorem 1.2. Let $\{\eta_t\}$ be a Lévy process. Then η_t has a càdlàg version (right continuous with left limits) which is also a Lévy process.

Proof. See, e.g., [P, S]. □

In view of this result we will from now on assume that the Lévy processes we work with are càdlàg.

The *jump* of η_t at $t \geq 0$ is defined by

$$\Delta\eta_t = \eta_t - \eta_{t-}. \quad (1.1.1)$$

Let \mathbf{B}_0 be the family of Borel sets $U \subset \mathbb{R}$ whose closure \bar{U} does not contain 0. For $U \in \mathbf{B}_0$ we define

$$N(t, U) = N(t, U, \omega) = \sum_{s: 0 < s \leq t} \mathcal{X}_U(\Delta\eta_s). \quad (1.1.2)$$

In other words, $N(t, U)$ is the number of jumps of size $\Delta\eta_s \in U$ which occur before or at time t . $N(t, U)$ is called the *Poisson random measure* (or *jump measure*) of $\eta(\cdot)$.

Remark 1.3. Note that $N(t, U)$ is *finite* for all $U \in \mathbf{B}_0$. To see this we proceed as follows: Define

$$T_1(\omega) = \inf\{t > 0; \eta_t \in U\}.$$

We claim that $T_1(\omega) > 0$ a.s. To prove this note that by right continuity of paths we have

$$\lim_{t \rightarrow 0^+} \eta(t) = \eta(0) = 0 \quad \text{a.s.}$$

Therefore, for all $\varepsilon > 0$ there exists $t(\varepsilon) > 0$ such that $|\eta(t)| < \varepsilon$ for all $t < t(\varepsilon)$. This implies that $\eta(t) \notin U$ for all $t < t(\varepsilon)$, if $\varepsilon < \text{dist}(0, U)$.

Next define inductively

$$T_{n+1}(\omega) = \inf\{t > T_n(\omega); \Delta\eta_t \in U\}.$$

Then by the above argument $T_{n+1} > T_n$ a.s. We claim that

$$T_n \rightarrow \infty \quad \text{as } n \rightarrow \infty, \text{ a.s.}$$

Assume not. Then $T_n \rightarrow T < \infty$. But then

$$\lim_{s \rightarrow T^-} \eta(s) \quad \text{cannot exist,}$$

contradicting the existence of left limits of the paths.

It is well known that Brownian motion $\{B(t)\}_{t \geq 0}$ has stationary and independent increments. Thus $B(t)$ is a Lévy process. Another important example is the following.

Example 1.4 (The Poisson Process). The Poisson process $\pi(t)$ of intensity $\lambda > 0$ is a Lévy process taking values in $\mathbb{N} \cup \{0\}$ and such that

$$P[\pi(t) = n] = \frac{(\lambda t)^n}{n!} e^{-\lambda t}; \quad n = 0, 1, 2, \dots$$

Theorem 1.5. [P, Theorem 1.35].

1. The set function $U \rightarrow N(t, U, \omega)$ defines a σ -finite measure on \mathbf{B}_0 for each fixed t, ω . The differential form of this measure is written $N(t, dz)$.
2. The set function $[a, b) \times U \rightarrow N(b, U, \omega) - N(a, U, \omega)$; $[a, b) \subset [0, \infty)$, $U \in \mathbf{B}_0$ defines a σ -finite measure for each fixed ω . The differential form of this measure is written $N(dt, dz)$.
3. The set function

$$\nu(U) = E[N(1, U)], \tag{1.1.3}$$

where $E = E_P$ denotes expectation with respect to P , also defines a σ -finite measure on \mathbf{B}_0 , called the Lévy measure of $\{\eta_t\}$.

4. Fix $U \in \mathbf{B}_0$. Then the process

$$\pi_U(t) := \pi_U(t, \omega) := N(t, U, \omega)$$

is a Poisson process of intensity $\lambda = \nu(U)$.

Example 1.6 (The Compound Poisson Process). Let $X(n)$, $n \in \mathbb{N}$ be a sequence of i.i.d. random variables taking values in \mathbb{R} with common distribution $\mu_{X(1)} = \mu_X$ and let $\pi(t)$ be a Poisson process of intensity λ , independent of all the $X(n)$ s.

The *compound Poisson process* $Y(t)$ is defined by

$$Y(t) = X(1) + \cdots + X(\pi(t)), \quad t \geq 0. \quad (1.1.4)$$

An increment of this process is given by

$$Y(s) - Y(t) = \sum_{k=\pi(t)+1}^{\pi(s)} X(k), \quad s > t.$$

This is independent of $X(1), \dots, X(\pi(t))$, and its distribution depends only on the difference $(s - t)$ and on the distribution of $X(1)$. Thus $Y(t)$ is a Lévy process.

To find the Lévy measure ν of $Y(t)$ note that if $U \in \mathbf{B}_0$ then

$$\begin{aligned} \nu(U) &= E[N(1, U)] = E \left[\sum_{s; 0 < s \leq 1} \mathcal{X}_U(\Delta Y(s)) \right] \\ &= E[(\text{number of jumps}) \times X_U(\text{jump})] = E[\pi(1)\mathcal{X}_U(X)] = \lambda\mu_X(U), \end{aligned}$$

by independence. We conclude that

$$\nu = \lambda\mu_X. \quad (1.1.5)$$

This shows that a Lévy process can be represented by a compound Poisson process if and only if its Lévy measure is finite. Note, however, that there are many interesting Lévy processes η_t with infinite Lévy measure ν , in fact even with $\int_{|z| < 1} |z| \nu(dz) = \infty$. See, e.g., [B]. In general, one can prove that for any fixed $R > 0$ the processes

$$M_t^{(k)} := \int_{1/k \leq |z| \leq R} z(N(t, dz) - t\nu(dz)), \quad k = 1, 2, \dots$$

are $L^2(P)$ -martingales and they converge in $L^2(P)$ to a martingale M_t denoted by

$$M_t = \int_{|z| \leq R} z(N(t, dz) - t\nu(dz)).$$

In fact we have

Theorem 1.7 (Itô–Lévy Decomposition [JS]). *Let $\{\eta_t\}$ be a Lévy process. Then η_t has the decomposition*

$$\eta_t = \alpha t + \sigma B(t) + \int_{|z| < R} z \tilde{N}(t, dz) + \int_{|z| \geq R} z N(t, dz), \quad (1.1.6)$$

for some constants $\alpha \in \mathbb{R}$, $\sigma \in \mathbb{R}$, $R \in [0, \infty]$. Here

$$\tilde{N}(dt, dz) = N(dt, dz) - \nu(dz)dt \quad (1.1.7)$$

is the compensated Poisson random measure of $\eta(\cdot)$, and $B(t)$ is a Brownian motion independent of $\tilde{N}(dt, dz)$. For each $A \in \mathbf{B}_0$ the process

$$M_t := \tilde{N}(t, A) \quad \text{is a martingale.} \quad (1.1.8)$$

If $\alpha = 0$ and $R = \infty$, we call η_t a Lévy martingale.

Theorem 1.8. *We can always choose $R = 1$. If*

$$E|\eta_t| < \infty \quad \text{for all } t \geq 0,$$

then

$$\int_{|z| \geq 1} |z| \nu(dz) < \infty$$

and hence we may choose $R = \infty$ and write

$$\eta_t = \alpha t + \sigma B(t) + \int_{\mathbb{R}} z \tilde{N}(t, dz).$$

(See [S, Theorem 25.3].)

Theorem 1.9 ([P]). *A Lévy process is a strong Markov process.*

Theorem 1.10 (The Lévy–Khintchine formula [P]). *Let $\{\eta_t\}$ be a Lévy process with Lévy measure ν . Then $\int_{\mathbb{R}} \min(1, z^2) \nu(dz) < \infty$ and*

$$E[e^{iu\eta_t}] = e^{t\psi(u)}, \quad u \in \mathbb{R}, \quad (1.1.9)$$

where

$$\psi(u) = -\frac{1}{2}\sigma^2 u^2 + i\alpha u + \int_{|z| < R} \{e^{iuz} - 1 - iuz\} \nu(dz) + \int_{|z| \geq R} (e^{iuz} - 1) \nu(dz). \quad (1.1.10)$$

Conversely, given constants α, σ^2 , and a measure ν on \mathbf{B}_0 s.t.

$$\int_{\mathbb{R}} \min(1, z^2) \nu(dz) < \infty,$$

there exists a Lévy process $\eta(t)$ (unique in law) such that (1.1.9–1.1.10) hold.

Note. It is possible that $\int_{|z| \leq R} |z| \nu(dz) = \infty$.

Theorem 1.11. [P, Corollary p. 48]. A Lévy process is a semimartingale.

Definition 1.12. [P]. Let \mathbf{D}_{ucp} denote the space of càdlàg adapted processes, equipped with the topology of uniform convergence on compacts in probability (ucp) : $H_n \rightarrow H$ ucp if for all $t > 0$ $\sup_{0 \leq s \leq t} |H_n(s) - H(s)| \rightarrow 0$ in probability ($A_n \rightarrow A$ in probability if for all $\varepsilon > 0$ there exists $n_\varepsilon \in \mathbb{N}$ such that $n \geq n_\varepsilon \Rightarrow \text{Prob.}(|A_n - A| > \varepsilon) < \varepsilon$).

Let \mathbf{L}_{ucp} denote the space of adapted càglàd processes (left continuous with right limits), equipped with the ucp topology. If $H(t)$ is a step function of the form

$$H(t) = H_0 \mathcal{X}_{\{0\}}(t) + \sum_i H_i \mathcal{X}_{(T_i, T_{i+1}]}(t),$$

where $H_i \in \mathcal{F}_{T_i}$ and $0 = T_0 \leq T_1 \leq \dots \leq T_{n+1} < \infty$ are \mathcal{F}_t -stopping times and X is càdlàg, we define

$$J_X H(t) := \int_0^t H_s dX_s := H_0 X_0 + \sum_i H_i (X_{T_{i+1} \wedge t} - X_{T_i \wedge t}), \quad t \geq 0.$$

Theorem 1.13. [P, p. 51]. Let X be a semimartingale. Then the mapping J_X can be extended to a continuous linear map

$$J_X : \mathbf{L}_{\text{ucp}} \rightarrow \mathbf{D}_{\text{ucp}}.$$

This construction allows us to define stochastic integrals of the form

$$\int_0^t H(s) d\eta_s$$

for all $H \in \mathbf{L}_{\text{ucp}}$. (See also Remark 1.18.) In view of the decomposition (1.1.6) this integral can be split into integrals with respect to ds , $dB(s)$, $\tilde{N}(ds, dz)$, and $N(ds, dz)$. This makes it natural to consider the more general stochastic integrals of the form

$$X(t) = X(0) + \int_0^t \alpha(s, \omega) ds + \int_0^t \beta(s, \omega) dB(s) + \int_0^t \int_{\mathbb{R}} \gamma(s, z, \omega) \tilde{N}(ds, dz), \tag{1.1.11}$$

where the integrands are satisfying the appropriate conditions for the integrals to exist and we for simplicity have put

$$\tilde{N}(ds, dz) = \begin{cases} N(ds, dz) - \nu(dz) ds & \text{if } |z| < R \\ N(ds, dz) & \text{if } |z| \geq R, \end{cases}$$

with R as in Theorem 1.7. As is customary we will use the following shorthand differential notation for processes $X(t)$ satisfying (1.1.11):

$$dX(t) = \alpha(t) dt + \beta(t) dB(t) + \int_{\mathbb{R}} \gamma(t, z) \tilde{N}(dt, dz). \tag{1.1.12}$$

We call such processes *Itô-Lévy processes*.

Recall that a semimartingale $M(t)$ is called a *local martingale* up to time T (with respect to P) if there exists an increasing sequence of \mathcal{F}_t -stopping times τ_n such that $\lim_{n \rightarrow \infty} \tau_n = T$ a.s. and

$M(t \wedge \tau_n)$ is a martingale with respect to P for all n .

Note that

1. If

$$E \left[\int_0^T \int_{\mathbb{R}} \gamma^2(t, z) \nu(dz) dt \right] < \infty, \quad (1.1.13)$$

then the process

$$M(t) := \int_0^t \int_{\mathbb{R}} \gamma(t, z) \tilde{N}(dt, dz), \quad 0 \leq t \leq T$$

is a *martingale*.

2. If

$$\int_0^T \int_{\mathbb{R}} \gamma^2(t, z) \nu(dz) dt < \infty \text{ a.s.}, \quad (1.1.14)$$

then $M(t)$ is a *local martingale*, $0 \leq t \leq T$.

1.2 The Itô Formula and Related Results

We now come to the important Itô formula for Itô–Lévy processes:

If $X(t)$ is given by (1.1.12) and $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ is a C^2 function, is the process $Y(t) := f(t, X(t))$ again an Itô–Lévy process and if so, how do we represent it in the form (1.1.12)?

If we argue heuristically and use our knowledge of the classical Itô formula it is easy to guess what the answer is:

Let $X^{(c)}(t)$ be the continuous part of $X(t)$, i.e., $X^{(c)}(t)$ is obtained by removing the jumps from $X(t)$. Then an increment in $Y(t)$ stems from an increment in $X^{(c)}(t)$ plus the jumps (coming from $N(\cdot, \cdot)$). Hence in view of the classical Itô formula we would guess that

$$\begin{aligned} dY(t) &= \frac{\partial f}{\partial t}(t, X(t))dt + \frac{\partial f}{\partial x}(t, X(t))dX^{(c)}(t) + \frac{1}{2} \frac{\partial^2 f}{\partial x^2}(t, X(t)) \cdot \beta^2(t)dt \\ &\quad + \int_{\mathbb{R}} \{f(t, X(t^-) + \gamma(t, z)) - f(t, X(t^-))\} N(dt, dz). \end{aligned}$$

It can be proved that our guess is correct. Since

$$dX^{(c)}(t) = \left(\alpha(t) - \int_{|z| < R} \gamma(t, z) \nu(dz) \right) dt + \beta(t)dB(t),$$

this gives the following result.

Theorem 1.14 (The One-Dimensional Itô Formula [BL, A, P]). *Suppose $X(t) \in \mathbb{R}$ is an Itô-Lévy process of the form*

$$dX(t) = \alpha(t, \omega)dt + \beta(t, \omega)dB(t) + \int_{\mathbb{R}} \gamma(t, z, \omega) \bar{N}(dt, dz), \quad (1.2.1)$$

where

$$\bar{N}(dt, dz) = \begin{cases} N(dt, dz) - \nu(dz)dt & \text{if } |z| < R \\ N(dt, dz) & \text{if } |z| \geq R \end{cases} \quad (1.2.2)$$

for some $R \in [0, \infty]$.

Let $f \in C^2(\mathbb{R}^2)$ and define $Y(t) = f(t, X(t))$. Then $Y(t)$ is again an Itô-Lévy process and

$$\begin{aligned} dY(t) &= \frac{\partial f}{\partial t}(t, X(t))dt + \frac{\partial f}{\partial x}(t, X(t))[\alpha(t, \omega)dt + \beta(t, \omega)dB(t)] \\ &\quad + \frac{1}{2}\beta^2(t, \omega) \frac{\partial^2 f}{\partial x^2}(t, X(t))dt \\ &\quad + \int_{|z| < R} \left\{ f(t, X(t^-) + \gamma(t, z)) - f(t, X(t^-)) \right. \\ &\quad \quad \left. - \frac{\partial f}{\partial x}(t, X(t^-))\gamma(t, z) \right\} \nu(dz)dt \\ &\quad + \int_{\mathbb{R}} \{f(t, X(t^-) + \gamma(t, z)) - f(t, X(t^-))\} \bar{N}(dt, dz). \end{aligned} \quad (1.2.3)$$

Note. If $R = 0$ then $\bar{N} = N$ everywhere.

If $R = \infty$ then $\bar{N} = \tilde{N}$ everywhere.

Example 1.15 (The Geometric Lévy Process). Consider the stochastic differential equation (SDE)

$$dX(t) = X(t^-) \left[\alpha dt + \beta dB(t) + \int_{\mathbb{R}} \gamma(t, z) \bar{N}(dt, dz) \right], \quad (1.2.4)$$

where α, β are constants and $\gamma(t, z) \geq -1$. To find the solution $X(t)$ of this equation we rewrite it as follows:

$$\frac{dX(t)}{X(t^-)} = \alpha dt + \beta dB(t) + \int_{\mathbb{R}} \gamma(t, z) \bar{N}(dt, dz).$$

Now define

$$Y(t) = \ln X(t).$$

Then by Itô's formula,

$$\begin{aligned}
dY(t) &= \frac{X(t)}{X(t)} [\alpha dt + \beta dB(t)] - \frac{1}{2} \beta^2 X^{-2}(t) X^2(t) dt \\
&\quad + \int_{|z| < R} \{ \ln(X(t^-) + \gamma(t, z) X(t^-)) - \ln(X(t^-)) \\
&\quad - X^{-1}(t^-) \gamma(t, z) X(t^-) \} \nu(dz) dt \\
&\quad + \int_{\mathbb{R}} \{ \ln(X(t^-) + \gamma(t, z) X(t^-)) - \ln(X(t^-)) \} \bar{N}(dt, dz) \\
&= \left(\alpha - \frac{1}{2} \beta^2 \right) dt + \beta dB(t) + \int_{|z| < R} \{ \ln(1 + \gamma(t, z)) - \gamma(t, z) \} \nu(dz) dt \\
&\quad + \int_{\mathbb{R}} \ln(1 + \gamma(t, z)) \bar{N}(dt, dz).
\end{aligned}$$

Hence

$$\begin{aligned}
Y(t) &= Y(0) + \left(\alpha - \frac{1}{2} \beta^2 \right) t + \beta B(t) + \int_0^t \int_{|z| < R} \{ \ln(1 + \gamma(s, z)) \\
&\quad - \gamma(s, z) \} \nu(dz) ds + \int_0^t \int_{\mathbb{R}} \ln(1 + \gamma(s, z)) \bar{N}(ds, dz)
\end{aligned}$$

and this gives the solution

$$\begin{aligned}
X(t) &= X(0) \exp \left\{ \left(\alpha - \frac{1}{2} \beta^2 \right) t + \beta B(t) \right. \\
&\quad + \int_0^t \int_{|z| < R} \{ \ln(1 + \gamma(s, z)) - \gamma(s, z) \} \nu(dz) ds \\
&\quad \left. + \int_0^t \int_{\mathbb{R}} \ln(1 + \gamma(s, z)) \bar{N}(ds, dz) \right\}. \quad (1.2.5)
\end{aligned}$$

In analogy with the diffusion case ($N = 0$) we call this process $X(t)$ a *geometric Lévy process*. It is often used as a model for stock prices. See, e.g., [B].

Next we formulate the corresponding multidimensional version of Theorem 1.14.

Theorem 1.16 (The Multidimensional Itô Formula). *Let $X(t) \in \mathbb{R}^n$ be an Itô-Lévy process of the form*

$$dX(t) = \alpha(t, \omega) dt + \sigma(t, \omega) dB(t) + \int_{\mathbb{R}^\ell} \gamma(t, z, \omega) \bar{N}(dt, dz), \quad (1.2.6)$$

where $\alpha : [0, T] \times \Omega \rightarrow \mathbb{R}^n$, $\sigma : [0, T] \times \Omega \rightarrow \mathbb{R}^{n \times m}$, and $\gamma : [0, T] \times \mathbb{R}^\ell \times \Omega \rightarrow \mathbb{R}^{n \times \ell}$ are adapted processes such that the integrals exist. Here $B(t)$ is an m -dimensional Brownian motion and

$$\begin{aligned}
 \bar{N}(dt, dz)^T &= (\bar{N}_1(dt, dz_1), \dots, \bar{N}_\ell(dt, dz_\ell)) \\
 &= (N_1(dt, dz_1) - \mathcal{X}_{|z_1| < R_1} \nu_1(dz_1)dt, \dots, N_\ell(dt, dz_\ell) \\
 &\quad - \mathcal{X}_{|z_\ell| < R_\ell} \nu_\ell(dz_\ell)dt),
 \end{aligned}$$

where $\{N_j\}$ are independent Poisson random measures with Lévy measures ν_j coming from ℓ independent (one-dimensional) Lévy processes η_1, \dots, η_ℓ .

Note that each column $\gamma^{(k)}$ of the $n \times \ell$ matrix $\gamma = [\gamma_{ij}]$ depends on z only through the k th coordinate z_k , i.e.,

$$\gamma^{(k)}(t, z, \omega) = \gamma^{(k)}(t, z_k, \omega), \quad z = (z_1, \dots, z_\ell) \in \mathbb{R}^\ell.$$

Thus the integral on the right of (1.2.6) is just a shorthand matrix notation. When written out in detail component number i of $X(t)$ in (1.2.6), $X_i(t)$ gets the form

$$\begin{aligned}
 dX_i(t) &= \alpha_i(t, \omega)dt + \sum_{j=1}^m \sigma_{ij}(t, \omega)dB_j(t) \\
 &\quad + \sum_{j=1}^{\ell} \int_{\mathbb{R}} \gamma_{ij}(t, z_j, \omega) \bar{N}_j(dt, dz_j), \quad 1 \leq i \leq n. \quad (1.2.7)
 \end{aligned}$$

Let $f \in C^{1,2}([0, T] \times \mathbb{R}^n; \mathbb{R})$. Put $Y(t) = f(t, X(t))$. Then

$$\begin{aligned}
 dY(t) &= \frac{\partial f}{\partial t} dt + \sum_{i=1}^n \frac{\partial f}{\partial x_i} (\alpha_i dt + \sigma_i dB(t)) + \frac{1}{2} \sum_{i,j=1}^n (\sigma \sigma^T)_{ij} \frac{\partial^2 f}{\partial x_i \partial x_j} dt \\
 &\quad + \sum_{k=1}^{\ell} \int_{|z_k| < R_k} \left\{ f(t, X(t^-) + \gamma^{(k)}(t, z_k)) - f(t, X(t^-)) \right. \\
 &\quad \quad \left. - \sum_{i=1}^n \gamma_i^{(k)}(t, z_k) \frac{\partial f}{\partial x_i}(X(t^-)) \right\} \nu_k(dz_k) dt \\
 &\quad + \sum_{k=1}^{\ell} \int_{\mathbb{R}} \{f(t, X(t^-) + \gamma^{(k)}(t, z_k)) - f(t, X(t^-))\} \bar{N}_k(dt, dz_k),
 \end{aligned} \quad (1.2.8)$$

where $\gamma^{(k)} \in \mathbb{R}^n$ is column number k of the $n \times \ell$ matrix $\gamma = [\gamma_{ik}]$ and $\gamma_i^{(k)} = \gamma_{ik}$ is the coordinate number i of $\gamma^{(k)}$.

Theorem 1.17 (The Itô–Lévy Isometry). Let $X(t) \in \mathbb{R}^n$ be as in (1.2.6) but with $X(0) = 0$ and $\alpha = 0$. Then

$$\begin{aligned}
E[X^2(T)] &= E \left[\int_0^T \left\{ \sum_{i=1}^n \sum_{j=1}^m \sigma_{ij}^2(t) + \sum_{i=1}^n \sum_{j=1}^{\ell} \int_{\mathbb{R}} \gamma_{ij}^2(t, z_j) \nu_j(dz_j) \right\} dt \right] \\
&= \sum_{i=1}^n E \left[\int_0^T \left\{ \sum_{j=1}^m \sigma_{ij}^2(t) + \sum_{j=1}^{\ell} \int_{\mathbb{R}} \gamma_{ij}^2(t, z_j) \nu_j(dz_j) \right\} dt \right] \quad (1.2.9)
\end{aligned}$$

provided that the right-hand side is finite.

Proof. This follows from the Itô formula applied to $f(t, x) = x^2 = |x|^2$. We omit the details. \square

Remark 1.18. As a special case of Theorem 1.17 assume that

$$dX(t) = d\eta(t) = \int_{\mathbb{R}} z \tilde{N}(dt, dz) \in \mathbb{R}$$

with $E[X^2(T)] = T \int_{\mathbb{R}} z^2 \nu(dz) < \infty$. Then we get the isometry

$$E \left[\left(\int_0^T H(t) d\eta(t) \right)^2 \right] = E \left[\int_0^T H^2(t) dt \right] \int_{\mathbb{R}} z^2 \nu(dz)$$

for all $H \in \mathbf{L}_{\text{ucp}}$ (see Definition 1.12) such that $H \in L^2([0, T] \times \Omega)$, i.e., such that

$$\|H\|_{L^2([0, T] \times \Omega)}^2 := E \left[\int_0^T H^2(t) dt \right] < \infty.$$

Using this we can in the usual way extend the definition of the integral

$$\int_0^T Y(t) d\eta(t) \in L^2(\Omega)$$

to all processes $Y(t)$ which are limits in $L^2([0, T] \times \Omega)$ of processes $H_n(t) \in \mathbf{L}_{\text{ucp}} \cap L^2([0, T] \times \Omega)$. We will call such processes $Y(t)$ *predictable processes*.

1.3 Lévy Stochastic Differential Equations

The geometric Lévy process is an example of a *Lévy diffusion*, i.e., the solution of a SDE driven by Lévy processes.

Theorem 1.19 (Existence and Uniqueness of Solutions of Lévy SDEs). *Consider the following Lévy SDE in \mathbb{R}^n : $X(0) = x_0 \in \mathbb{R}^n$ and*

$$dX(t) = \alpha(t, X(t))dt + \sigma(t, X(t))dB(t) + \int_{\mathbb{R}^n} \gamma(t, X(t^-), z) \tilde{N}(dt, dz), \quad (1.3.1)$$

where $\alpha : [0, T] \times \mathbb{R}^n \rightarrow \mathbb{R}^n$, $\sigma : [0, T] \times \mathbb{R}^n \rightarrow \mathbb{R}^{n \times m}$ and $\gamma : [0, T] \times \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}^{n \times \ell}$ satisfy the following conditions

(At most linear growth) There exists a constant $C_1 < \infty$ such that

$$\|\sigma(t, x)\|^2 + |\alpha(t, x)|^2 + \int_{\mathbb{R}} \sum_{k=1}^{\ell} |\gamma_k(t, x, z)|^2 \nu_k(dz_k) \leq C_1(1 + |x|^2)$$

for all $x \in \mathbb{R}^n$

(Lipschitz continuity) There exists a constant $C_2 < \infty$ such that

$$\begin{aligned} & \|\sigma(t, x) - \sigma(t, y)\|^2 + |\alpha(t, x) - \alpha(t, y)|^2 \\ & + \sum_{k=1}^{\ell} \int_{\mathbb{R}} |\gamma^{(k)}(t, x, z_k) - \gamma^{(k)}(t, y, z_k)|^2 \nu_k(dz_k) \leq C_2|x - y|^2, \end{aligned}$$

for all $x, y \in \mathbb{R}^n$.

Then there exists a unique càdlàg adapted solution $X(t)$ such that

$$E[|X(t)|^2] < \infty \quad \text{for all } t.$$

Solutions of Lévy SDEs in the *time homogeneous* case, i.e., when $\alpha(t, x) = \alpha(x)$, $\sigma(t, x) = \sigma(x)$, and $\gamma(t, x, z) = \gamma(x, z)$ are called *jump diffusions* (or *Lévy diffusions*).

Theorem 1.20. *A jump diffusion is a strong Markov process.*

Proof. See [P, Theorem V.32]. □

Definition 1.21. *Let $X(t) \in \mathbb{R}^n$ be a jump diffusion. Then the generator A of X is defined on functions $f : \mathbb{R}^n \rightarrow \mathbb{R}$ by*

$$Af(x) = \lim_{t \rightarrow 0^+} \frac{1}{t} \{E^x[f(X(t))] - f(x)\} \quad (\text{if the limit exists}),$$

where $E^x[f(X(t))] = E[f(X^{(x)}(t))]$, $X^{(x)}(0) = x$.

Theorem 1.22. *Suppose $f \in C_0^2(\mathbb{R}^n)$. Then $Af(x)$ exists and is given by*

$$\begin{aligned} Af(x) &= \sum_{i=1}^n \alpha_i(x) \frac{\partial f}{\partial x_i}(x) + \frac{1}{2} \sum_{i,j=1}^n (\sigma \sigma^T)_{ij}(x) \frac{\partial^2 f}{\partial x_i \partial x_j}(x) \\ &+ \int_{\mathbb{R}} \sum_{k=1}^{\ell} \{f(x + \gamma^{(k)}(x, z)) - f(x) - \nabla f(x) \cdot \gamma^{(k)}(x, z)\} \nu_k(dz_k). \end{aligned} \tag{1.3.2}$$

From now on we *define* $Af(x)$ by the expression (1.3.2) for all f such that the partial derivatives of f and the integrals in (1.3.2) exist at x .

Theorem 1.23 (The Dynkin Formula I). *Let $X(t) \in \mathbb{R}^n$ be a jump diffusion and let $f \in C_0^2(\mathbb{R}^n)$. Let τ be a stopping time such that*

$$E^x[\tau] < \infty.$$

Then

$$E^x[f(X(\tau))] = f(x) + E^x \left[\int_0^\tau Af(X(s))ds \right].$$

Proof. This follows by combining the Itô formula (1.2.8) with the formula (1.3.2) for A and taking expectation. \square

This version is usually strong enough for applications in the case when there are no jumps ($N = 0$). However, for jump diffusions we need the following stronger, localized version.

Theorem 1.24 (The Dynkin Formula II). *Let $X(t) \in \mathbb{R}^n$ be a jump diffusion, $G \subset \mathbb{R}^n$ be an open set and let $f \in C^2(G) \cap C(\bar{G})$. Let $\tau < \infty$ be a stopping time. Suppose that*

$$\tau \leq \tau_G := \inf\{t > 0; X(t) \notin G\} \quad (1.3.3)$$

$$X(\tau) \in \bar{G} \quad \text{a.s.} \quad (1.3.4)$$

$$E^x \left[|f(X(\tau))| + \int_0^\tau |Af(X(t))|dt \right] < \infty. \quad (1.3.5)$$

Then we have

$$E^x[f(X(\tau))] = f(x) + E^x \left[\int_0^\tau (Af)(X(t))dt \right].$$

Definition 1.25. *In general, if $\{\psi_m\}_{m=1}^\infty$ and g are functions defined on a set $G \subset \mathbb{R}^n$, we say that $\psi_m \rightarrow g$ pointwise dominatedly in G if $\psi_m(x) \rightarrow g(x)$ for all $x \in G$ and there exists a constant $C < \infty$ such that*

$$|\psi_m(x)| \leq C|g(x)| \quad \text{for all } x \in G, m = 1, 2, \dots$$

Proof of Theorem 1.24. Choose $f_m \in C_0^2(\mathbb{R}^n)$ such that $f_m \rightarrow f$ pointwise dominatedly in \bar{G} and $\frac{\partial f_m}{\partial x_i} \rightarrow \frac{\partial f}{\partial x_i}$, $\frac{\partial^2 f_m}{\partial x_i \partial x_j} \rightarrow \frac{\partial^2 f}{\partial x_i \partial x_j}$, and $Af_m \rightarrow Af$ pointwise dominatedly in G for all $i, j = 1, \dots, n$. Then apply Theorem 1.23 to each f_m and $\tau \wedge k$, $k = 1, 2, \dots$. Let $m, k \rightarrow \infty$ and apply the dominated convergence theorem. \square

1.4 The Girsanov Theorem and Applications

The Girsanov theorem and the related concept of an equivalent local martingale measure (ELMM) are important in the applications of stochastic analysis to finance. In this chapter we first give a general semimartingale discussion

and then we apply it to Itô–Lévy processes. We refer to [Ka] for more details.

Let $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \geq 0}, P)$ be a filtered probability space. Let Q be another probability measure on \mathcal{F}_T . We say that Q is *equivalent* to $P \mid \mathcal{F}_T$ if $P \mid \mathcal{F}_T \ll Q$ and $Q \ll P \mid \mathcal{F}_T$, or, equivalently, if P and Q have the same zero sets in \mathcal{F}_T . By the Radon–Nikodym theorem this is the case if and only if we have

$$dQ(\omega) = Z(T)dP(\omega) \quad \text{and} \quad dP(\omega) = Z^{-1}(T)dQ(\omega) \quad \text{on } \mathcal{F}_T$$

for some \mathcal{F}_T -measurable random variable $Z(T) > 0$ a.s. P . In that case we also write

$$\frac{dQ}{dP} = Z(T) \quad \text{and} \quad \frac{dP}{dQ} = Z^{-1}(T) \quad \text{on } \mathcal{F}_T. \quad (1.4.1)$$

We first make a simple, but useful observation.

Lemma 1.26. *Suppose $Q \ll P$ with $\frac{dQ}{dP} = Z(T)$ on \mathcal{F}_T . Then*

$$Q \mid \mathcal{F}_t \ll P \mid \mathcal{F}_t \quad \text{for all } t \in [0, T] \quad \text{and}$$

$$Z(t) := \frac{d(Q \mid \mathcal{F}_t)}{d(P \mid \mathcal{F}_t)} = E_P[Z(T) \mid \mathcal{F}_t], \quad 0 \leq t \leq T. \quad (1.4.2)$$

In particular, $Z(t)$ is a P -martingale.

Proof of Theorem 1.24. Since $P(G) = 0 \Rightarrow Q(G) = 0$ for all $G \in \mathcal{F}_T \supseteq \mathcal{F}_t$, it is clear that $Q \mid \mathcal{F}_t \ll P \mid \mathcal{F}_t$. Choose $F \in \mathcal{F}_t$. Then

$$\begin{aligned} E_P[F \cdot E[Z(T) \mid \mathcal{F}_t]] &= E_P[E_P[FZ(T) \mid \mathcal{F}_t]] \\ &= E_P[FZ(T)] = E_Q[F] = E_P[F \cdot Z(t)]. \end{aligned}$$

Since this holds for all $F \in \mathcal{F}_t$ we conclude that

$$E_P[Z(T) \mid \mathcal{F}_t] = Z(t), \quad \text{as claimed.} \quad \square$$

Lemma 1.27. *Suppose $Q \ll P$ with $\frac{dQ}{dP} = Z(T) > 0$ on \mathcal{F}_T . Let $X(t)$ be an adapted process such that $Z(t)X(t)$ is a martingale with respect to P . Then $X(t)$ is a martingale with respect to Q . Similarly, if $Z(t)X(t)$ is a local martingale with respect to P , then $X(t)$ is a local martingale with respect to Q .*

Proof of Theorem 1.24. We prove the last statement. Let $\tau \geq t$ be a stopping time, $\tau \leq T$. Then

$$\begin{aligned} E_Q[X(\tau) \mid \mathcal{F}_t] &= \frac{E[Z(T)X(\tau) \mid \mathcal{F}_t]}{E[Z(T) \mid \mathcal{F}_t]} = \frac{E[Z(\tau)X(\tau) \mid \mathcal{F}_t]}{Z(t)} = \frac{Z(t)X(t)}{Z(t)} \\ &= X(t) \quad \text{for all } t \in [0, T]. \quad \square \end{aligned}$$

- [download The Rise and Fall of Classical Greece \(The Princeton History of the Ancient World Series\) here](#)
- [**Buzzed: The Straight Facts About the Most Used and Abused Drugs from Alcohol to Ecstasy \(Fully Revised and Updated 4th Edition\) pdf, azw \(kindle\), epub**](#)
- [The Thirteenth Step: Addiction in the Age of Brain Science pdf](#)
- [download Santorini Volcano \(Geological Society Memoir, Volume 19\)](#)
- [read online Community: Seeking Safety in an Insecure World pdf, azw \(kindle\), epub, doc, mobi](#)
- [The Order of Good Cheer pdf, azw \(kindle\), epub](#)

- <http://korplast.gr/lib/The-Rise-and-Fall-of-Classical-Greece--The-Princeton-History-of-the-Ancient-World-Series-.pdf>
- <http://betsy.wesleychapelcomputerrepair.com/library/Buzzed--The-Straight-Facts-About-the-Most-Used-and-Abused-Drugs-from-Alcohol-to-Ecstasy--Fully-Revised-and-Upda>
- <http://schrolf.de/books/The-Thirteenth-Step--Addiction-in-the-Age-of-Brain-Science.pdf>
- <http://academialanguagebar.com/?ebooks/Raising-Happy-Children-For-Dummies.pdf>
- <http://cavalldecartro.highlandagency.es/library/The-Barsoom-Project--Dream-Park--Book-2-.pdf>
- <http://pittiger.com/lib/The-Order-of-Good-Cheer.pdf>