

---

# **N. BOURBAKI**

## **ELEMENTS OF MATHEMATICS**

### **Functions of a Real Variable**

**Elementary Theory**



**Springer**

---

# ELEMENTS OF MATHEMATICS

---

**Springer**

*Berlin*

*Heidelberg*

*New York*

*Hong Kong*

*London*

*Milan*

*Paris*

*Tokyo*

---

NICOLAS BOURBAKI

ELEMENTS OF MATHEMATICS

Functions of a Real Variable

Elementary Theory



Springer

---

Originally published as  
Fonctions d'une variable réelle  
©Hermann, Paris, 1976 and Nicolas Bourbaki, 1982

Translator  
Philip Spain  
University of Glasgow  
Department of Mathematics  
University Gardens  
Glasgow G12 8QW  
Scotland  
e-mail: pgs@maths.gla.ac.uk

This work has been published with the help  
of the French *Ministère de la Culture and Centre national du livre*

Mathematics Subject Classification (2000): 26-02, 26A06, 26A12,  
26A15, 26A24, 26A42, 26A51, 34A12, 34A30, 46B03

Cataloging-in-Publication Data applied for  
A catalog record for this book is available from the Library of Congress.  
Bibliographic information published by Die Deutsche Bibliothek  
Die Deutsche Bibliothek lists this publication in the Deutsche Nationalbibliografie;  
detailed bibliographic data is available in the Internet at <http://dnb.ddb.de>

ISBN-13:978-3-642-63932-6 e-ISBN-13:978-3-642-59315-4  
DOI: 10.1007/978-3-642-59315-4

**Springer-Verlag Berlin Heidelberg New York**

This work is subject to copyright. All rights are reserved, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilm or in any other way, and storage in data banks. Duplication of this publication or parts thereof is permitted only under the provisions of the German Copyright Law of September 9, 1965, in its current version, and permission for use must always be obtained from Springer-Verlag. Violations are liable for prosecution under the German Copyright Law.

Springer-Verlag Berlin Heidelberg New York  
a member of BertelsmannSpringer Science+Business Media GmbH  
<http://www.springer.de>

© Springer-Verlag Berlin Heidelberg 2004  
Softcover reprint of the hardcover 1st edition 2004

The use of general descriptive names, registered names, trademarks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

Typesetting: by the Translator and Frank Herweg, Leutershausen  
Cover Design: *Design & Production* GmbH, Heidelberg  
Printed on acid-free paper 41/3142 db 5 4 3 2 1 0

---

## To the reader

1. The Elements of Mathematics Series takes up mathematics at the beginning, and gives complete proofs. In principle, it requires no particular knowledge of mathematics on the readers' part, but only a certain familiarity with mathematical reasoning and a certain capacity for abstract thought. Nevertheless it is directed especially to those who have a good knowledge of at least the content of the first year or two of a university mathematics course.

2. The method of exposition we have chosen is axiomatic, and normally proceeds from the general to the particular. The demands of proof impose a rigorously fixed order on the subject matter. It follows that the utility of certain considerations may not be immediately apparent to the reader until later chapters unless he has already a fairly extended knowledge of mathematics.

3. The series is divided into Books and each Book into chapters. The Books already published, either in whole or in part, in the French edition, are listed below. When an English translation is available, the corresponding English title is mentioned between parentheses. Throughout the volume a reference indicates the English edition, when available, and the French edition otherwise.

Théorie des Ensembles (Theory of Sets)	designated by	E	( <i>Set Theory</i> )
Algèbre (Algebra) <sup>1</sup>	—	A	( <i>Alg</i> )
Topologie Générale (General Topology)	—	TG	( <i>Gen. Top.</i> )
Fonctions d'une Variable Réelle (Functions of a Real Variable) <sup>2</sup>	—	FVR	( <i>FRV</i> )
Espaces Vectoriels Topologiques (Topological Vector Spaces)	—	EVT	( <i>Top. Vect. Sp.</i> )
Intégration	—	INT	
Algèbre Commutative (Commutative Algebra) <sup>3</sup>	—	AC	( <i>Comm. Alg.</i> )
Variétés Différentielles et Analytiques	—	VAR	
Groupes et Algèbres de Lie (Lie Groups and Lie Algebras) <sup>4</sup>	—	LIE	( <i>LIE</i> )
Théories Spectrales		TS	

In the first six Books (according to the above order), every statement in the text assumes as known only those results which have already discussed in the same

---

<sup>1</sup> So far, chapters I to VII only have been translated.

<sup>2</sup> This volume!

<sup>3</sup> So far, chapters I to VII only have been translated.

<sup>4</sup> So far, chapters I to III only have been translated.

chapter, or in the previous chapters ordered as follows: E ; A, chapters I to III ; TG, chapters I to III ; A, from chapter IV on ; TG, from chapter IV on ; FVR ; EVT ; INT.

From the seventh Book on, the reader will usually find a precise indication of its logical relationship to the other Books (the first six Books being always assumed to be known).

4. However, we have sometimes inserted examples in the text which refer to facts which the reader may already know but which have not yet been discussed in the Series. Such examples are placed between two asterisks : \* . . . \*. Most readers will undoubtedly find that these examples will help them to understand the text. In other cases, the passages between \* . . . \* refer to results which are discussed elsewhere in the Series. We hope the reader will be able to verify the absence of any vicious circle.

5. The logical framework of each chapter consists of the *definitions*, the *axioms*, and the *theorems* of the chapter. These are the parts that have mainly to be borne in mind for subsequent use. Less important results and those which can easily be deduced from the theorems are labelled as “propositions”, “lemmas”, “corollaries”, “remarks”, etc. Those which may be omitted at a first reading are printed in small type. A commentary on a particularly important theorem appears occasionally under the name of “scholium”.

To avoid tedious repetitions it is sometimes convenient to introduce notation or abbreviations which are in force only within a certain chapter or a certain section of a chapter (for example, in a chapter which is concerned only with commutative rings, the word “ring” would always signify “commutative ring”). Such conventions are always explicitly mentioned, generally at the beginning of the *chapter* in which they occur.

6. Some passages are designed to forewarn the reader against serious errors.

These passages are signposted in the margin with the sign (“dangerous bend”).

7. The Exercises are designed both to enable the reader to satisfy himself that he has digested the text and to bring to his notice results which have no place in the text but which are nonetheless of interest. The most difficult exercises bear the sign ¶.

8. In general we have adhered to the commonly accepted terminology, *except where there appeared to be good reasons for deviating from it*.

9. We have made a particular effort always to use rigorously correct language, without sacrificing simplicity. As far as possible we have drawn attention in the text to *abuses of language*, without which any mathematical text runs the risk of pedantry, not to say unreadability.

10. Since in principle the text consists of a dogmatic exposition of a theory, it contains in general no references to the literature. Bibliographical are gathered together in *Historical Notes*. The bibliography which follows each historical note contains in general only those books and original memoirs which have been of the greatest importance in the evolution of the theory under discussion. It makes no sort of pretence to completeness.

As to the exercises, we have not thought it worthwhile in general to indicate their origins, since they have been taken from many different sources (original papers, textbooks, collections of exercises).

11. In the present Book, references to theorems, axioms, definitions, . . . are given by quoting successively:

- the Book (using the abbreviation listed in Section 3), chapter and page, where they can be found ;
- the chapter and page only when referring to the present Book.

The *Summaries of Results* are quoted by to the letter R; thus *Set Theory*, R signifies “*Summary of Results of the Theory of Sets*”.



---

# CONTENTS

<b>TO THE READER</b> .....	V
<b>CONTENTS</b> .....	IX

---

## CHAPTER I DERIVATIVES

---

<b>§ 1. FIRST DERIVATIVE</b> .....	3
1. Derivative of a vector function .....	3
2. Linearity of differentiation .....	5
3. Derivative of a product .....	6
4. Derivative of the inverse of a function .....	8
5. Derivative of a composite function .....	9
6. Derivative of an inverse function .....	9
7. Derivatives of real-valued functions .....	10
<b>§ 2. THE MEAN VALUE THEOREM</b> .....	12
1. Rolle's Theorem .....	12
2. The mean value theorem for real-valued functions .....	13
3. The mean value theorem for vector functions .....	15
4. Continuity of derivatives .....	18
<b>§ 3. DERIVATIVES OF HIGHER ORDER</b> .....	20
1. Derivatives of order $n$ .....	20
2. Taylor's formula .....	21
<b>§ 4. CONVEX FUNCTIONS OF A REAL VARIABLE</b> .....	23
1. Definition of a convex function .....	24
2. Families of convex functions .....	27
3. Continuity and differentiability of convex functions .....	27
4. Criteria for convexity .....	30
Exercises on §1 .....	35
Exercises on §2 .....	37

Exercises on §3 .....	39
Exercises on §4 .....	45

---

## CHAPTER II PRIMITIVES AND INTEGRALS

---

<b>§ 1. PRIMITIVES AND INTEGRALS</b> .....	51
1. Definition of primitives .....	51
2. Existence of primitives .....	52
3. Regulated functions .....	53
4. Integrals .....	56
5. Properties of integrals .....	59
6. Integral formula for the remainder in Taylor's formula; primitives of higher order .....	62
<b>§ 2. INTEGRALS OVER NON-COMPACT INTERVALS</b> .....	62
1. Definition of an integral over a non-compact interval .....	62
2. Integrals of positive functions over a non-compact interval .....	66
3. Absolutely convergent integrals .....	67
<b>§ 3. DERIVATIVES AND INTEGRALS OF FUNCTIONS DEPENDING ON A PARAMETER</b> .....	68
1. Integral of a limit of functions on a compact interval .....	68
2. Integral of a limit of functions on a non-compact interval .....	69
3. Normally convergent integrals .....	72
4. Derivative with respect to a parameter of an integral over a compact interval .....	73
5. Derivative with respect to a parameter of an integral over a non-compact interval .....	75
6. Change of order of integration .....	76
Exercises on §1 .....	79
Exercises on §2 .....	84
Exercises on §3 .....	86

---

## CHAPTER III ELEMENTARY FUNCTIONS

---

<b>§ 1. DERIVATIVES OF THE EXPONENTIAL AND CIRCULAR FUNCTIONS</b> .....	91
1. Derivatives of the exponential functions; the number $e$ .....	91
2. Derivative of $\log_a x$ .....	93
3. Derivatives of the circular functions; the number $\pi$ .....	94
4. Inverse circular functions .....	95

5. The complex exponential .....	97
6. Properties of the function $e^z$ .....	98
7. The complex logarithm .....	100
8. Primitives of rational functions .....	101
9. Complex circular functions; hyperbolic functions .....	102
<b>§ 2. EXPANSIONS OF THE EXPONENTIAL AND CIRCULAR FUNCTIONS, AND OF THE FUNCTIONS ASSOCIATED WITH THEM</b> .....	105
1. Expansion of the real exponential .....	105
2. Expansions of the complex exponential, of $\cos x$ and $\sin x$ . .....	106
3. The binomial expansion .....	107
4. Expansions of $\log(1+x)$ , of $\text{Arc tan } x$ and of $\text{Arc sin } x$ .....	111
Exercises on §1 .....	115
Exercises on §2 .....	125
Historical Note (Chapters I-II-III) .....	129
Bibliography .....	159

---

## CHAPTER IV DIFFERENTIAL EQUATIONS

---

<b>§ 1. EXISTENCE THEOREMS</b> .....	163
1. The concept of a differential equation .....	163
2. Differential equations admitting solutions that are primitives of regulated functions .....	164
3. Existence of approximate solutions .....	166
4. Comparison of approximate solutions .....	168
5. Existence and uniqueness of solutions of Lipschitz and locally Lipschitz equations .....	171
6. Continuity of integrals as functions of a parameter .....	174
7. Dependence on initial conditions .....	176
<b>§ 2. LINEAR DIFFERENTIAL EQUATIONS</b> .....	177
1. Existence of integrals of a linear differential equation .....	177
2. Linearity of the integrals of a linear differential equation .....	179
3. Integrating the inhomogeneous linear equation .....	182
4. Fundamental systems of integrals of a linear system of scalar differential equations .....	183
5. Adjoint equation .....	186
6. Linear differential equations with constant coefficients .....	188
7. Linear equations of order $n$ .....	192
8. Linear equations of order $n$ with constant coefficients .....	194
9. Systems of linear equations with constant coefficients .....	196

Exercises on §1 .....	199
Exercises on §2 .....	204
Historical Note .....	207
Bibliography .....	209

---

## CHAPTER V LOCAL STUDY OF FUNCTIONS

---

<b>§ 1. COMPARISON OF FUNCTIONS ON A FILTERED SET</b> .....	211
1. Comparison relations: I. Weak relations .....	211
2. Comparison relations: II. Strong relations .....	214
3. Change of variable .....	217
4. Comparison relations between strictly positive functions .....	217
5. Notation .....	219
<b>§ 2. ASYMPTOTIC EXPANSIONS</b> .....	220
1. Scales of comparison .....	220
2. Principal parts and asymptotic expansions .....	221
3. Sums and products of asymptotic expansions .....	223
4. Composition of asymptotic expansions .....	224
5. Asymptotic expansions with variable coefficients .....	226
<b>§ 3. ASYMPTOTIC EXPANSIONS OF FUNCTIONS OF A REAL VARIABLE</b> .....	227
1. Integration of comparison relations: I. Weak relations .....	228
2. Application: logarithmic criteria for convergence of integrals .....	229
3. Integration of comparison relations: II. Strong relations .....	230
4. Differentiation of comparison relations .....	232
5. Principal part of a primitive .....	233
6. Asymptotic expansion of a primitive .....	235
<b>§ 4. APPLICATION TO SERIES WITH POSITIVE TERMS</b> .....	236
1. Convergence criteria for series with positive terms .....	236
2. Asymptotic expansion of the partial sums of a series .....	238
3. Asymptotic expansion of the partial products of an infinite product .....	243
4. Application: convergence criteria of the second kind for series with positive terms .....	244
<b>APPENDIX</b> .....	247
1. Hardy fields .....	247
2. Extension of a Hardy field .....	248
3. Comparison of functions in a Hardy field .....	250

4. (H) Functions .....	252
5. Exponentials and iterated logarithms .....	253
6. Inverse function of an (H) function .....	255
Exercises on §1 .....	259
Exercises on §3 .....	260
Exercises on §4 .....	261
Exercises on Appendix .....	263

---

**CHAPTER VI GENERALIZED TAYLOR EXPANSIONS.  
EULER-MACLAURIN SUMMATION FORMULA**

---

<b>§ 1. GENERALIZED TAYLOR EXPANSIONS</b> .....	269
1. Composition operators on an algebra of polynomials .....	269
2. Appell polynomials attached to a composition operator .....	272
3. Generating series for the Appell polynomials .....	274
4. Bernoulli polynomials .....	275
5. Composition operators on functions of a real variable .....	277
6. Indicatrix of a composition operator .....	278
7. The Euler-Maclaurin summation formula .....	282
<b>§ 2. EULERIAN EXPANSIONS OF THE TRIGONOMETRIC FUNCTIONS AND BERNOULLI NUMBERS</b> .....	283
1. Eulerian expansion of $\cot z$ .....	283
2. Eulerian expansion of $\sin z$ .....	286
3. Application to the Bernoulli numbers .....	287
<b>§ 3. BOUNDS FOR THE REMAINDER IN THE EULER-MACLAURIN SUMMATION FORMULA</b> .....	288
1. Bounds for the remainder in the Euler-Maclaurin summation formula .....	288
2. Application to asymptotic expansions .....	289
Exercises on §1 .....	291
Exercises on §2 .....	292
Exercises on §3 .....	296
Historical Note (Chapters V and VI) .....	299
Bibliography .....	303

---

**CHAPTER VII THE GAMMA FUNCTION**

---

<b>§ 1. THE GAMMA FUNCTION IN THE REAL DOMAIN</b> .....	305
1. Definition of the Gamma function .....	305
2. Properties of the Gamma function .....	307
3. The Euler integrals .....	310
<b>§ 2. THE GAMMA FUNCTION IN THE COMPLEX DOMAIN</b> .....	315
1. Extending the Gamma function to $\mathbf{C}$ .....	315
2. The complements' relation and the Legendre-Gauss multiplication formula .....	316
3. Stirling's expansion .....	319
Exercises on §1 .....	325
Exercises on §2 .....	327
Historical Note .....	329
Bibliography .....	331
<b>INDEX OF NOTATION</b> .....	333
<b>INDEX</b> .....	335

---

# INTRODUCTION

The purpose of this Book is the elementary study of the infinitesimal properties of *one* real variable; the extension of these properties to functions of *several* real variables, or, all the more, to functions defined on more general spaces, will be treated only in later Books.

The results which we shall demonstrate will be useful above all in relation to (finite) real-valued functions of a real variable; but most of them extend without further argument to functions of a real variable taking values in a *topological vector space* over  $\mathbf{R}$  (see below); as these functions occur frequently in Analysis we shall state for them all results which are not specific to real-valued functions.

The notion of a topological vector space, of which we have just spoken, is defined and studied in detail in Book V of this Series; but we do not need *any* of the results of Book V in this Book; some definitions, however, are needed, and we shall reproduce them below for the convenience of the reader.

We shall not repeat the definition of a *vector space* over a (*commutative*) *field*  $\mathbf{K}$  (*Alg.*, II, p. 193).<sup>1</sup> A *topological vector space*  $E$  over a *topological field*  $\mathbf{K}$  is a vector space over  $\mathbf{K}$  endowed with a topology such that the functions  $\mathbf{x} + \mathbf{y}$  and  $\mathbf{x}t$  are *continuous* on  $E \times E$  and  $E \times \mathbf{K}$  respectively; in particular, such a topology is compatible with the structure of the additive group of  $E$ . All topological vector spaces considered in this Book are implicitly assumed to be Hausdorff. When the topological group  $E$  is complete one says that the topological vector space  $E$  is *complete*. Every *normed* vector space over a *valued field*  $\mathbf{K}$  (*Gen. Top.*, IX, p. 169)<sup>2</sup> is a topological vector space over  $\mathbf{K}$ .

Let  $E$  be a vector space (with or without a topology) over the real field  $\mathbf{R}$ ; if  $\mathbf{x}, \mathbf{y}$  are arbitrary points in  $E$  the set of points  $\mathbf{x}t + \mathbf{y}(1 - t)$  where  $t$  runs through the closed

---

<sup>1</sup> The elements (or *vectors*) of a vector space  $E$  over a commutative field  $\mathbf{K}$  will usually be denoted in this chapter by thick minuscules, and scalars by roman minuscules; most often we shall place the scalar  $t$  to the *right* in the product of a vector  $\mathbf{x}$  by  $t$ , writing the product as  $\mathbf{x}t$ ; on occasion we will allow ourselves to use the left notation  $t\mathbf{x}$  in certain cases where it is more convenient; also, sometimes we shall write the product of the scalar  $1/t$  ( $t \neq 0$ ) and the vector  $\mathbf{x}$  in the form  $\mathbf{x}/t$ .

<sup>2</sup> We recall that a *norm* on  $E$  is a real function  $\|\mathbf{x}\|$  defined on  $E$ , taking finite non-negative values, such that the relation  $\|\mathbf{x}\| = 0$  is equivalent to  $\mathbf{x} = 0$  and such that

$$\|\mathbf{x} + \mathbf{y}\| \leq \|\mathbf{x}\| + \|\mathbf{y}\| \quad \text{and} \quad \|\mathbf{x}t\| = \|\mathbf{x}\| \cdot |t|$$

for all  $t \in \mathbf{K}$  ( $|t|$  being the absolute value of  $t$  in  $\mathbf{K}$ ).

segment  $[0, 1]$  of  $\mathbf{R}$  is called the *closed segment* with endpoints  $\mathbf{x}, \mathbf{y}$ . One says that a subset  $A$  of  $E$  is *convex* if for any  $\mathbf{x}, \mathbf{y}$  in  $A$  the closed segment with endpoints  $\mathbf{x}$  and  $\mathbf{y}$  is contained in  $A$ . For example, an affine linear variety is convex; so is any closed segment; in  $\mathbf{R}^n$  any parallelotope (*Gen. Top.*, VI, p. 34) is convex. Every intersection of convex sets is convex.

We say that a topological vector space  $E$  over the field  $\mathbf{R}$  is *locally convex* if the origin (and thus any point of  $E$ ) has a fundamental system of *convex* neighbourhoods. Every *normed* space is locally convex; indeed, the balls  $\|\mathbf{x}\| \leq r$  ( $r > 0$ ) form a fundamental system of neighbourhoods of 0 in  $E$ , and each of these is convex, for the relations  $\|\mathbf{x}\| \leq r, \|\mathbf{y}\| \leq r$  imply that

$$\|\mathbf{x}t + \mathbf{y}(1 - t)\| \leq \|\mathbf{x}\|t + \|\mathbf{y}\|(1 - t) \leq r$$

for  $0 \leq t \leq 1$ .

Finally, a *topological algebra*  $A$  over a (commutative) *topological field*  $\mathbf{K}$  is an algebra over  $\mathbf{K}$  endowed with a topology for which the functions  $\mathbf{x} + \mathbf{y}$ ,  $\mathbf{xy}$  and  $\mathbf{xt}$  are continuous on  $A \times A$ ,  $A \times A$  and  $A \times \mathbf{K}$  respectively; when one endows  $A$  only with its topology and vector space structure over  $\mathbf{K}$  then  $A$  is a topological vector space. Every *normed algebra* over a *valued field*  $\mathbf{K}$  (*Gen. Top.*, IX, p. 175) is a topological algebra over  $\mathbf{K}$ .



---

# CHAPTER I

## Derivatives

### § 1. FIRST DERIVATIVE

As was said in the Introduction, in this chapter and the next we shall study the infinitesimal properties of functions which are defined on a subset of the real field  $\mathbf{R}$  and take their values in a *Hausdorff topological vector space*  $E$  over the field  $\mathbf{R}$ ; for brevity we shall say that such a function is a *vector function of a real variable*. The most important case is that where  $E = \mathbf{R}$  (real-valued functions of a real variable). When  $E = \mathbf{R}^n$ , consideration of a vector function with values in  $E$  reduces to the simultaneous consideration of  $n$  finite real functions.

Many of the definitions and properties stated in chapter I extend to functions which are defined on a subset of the field  $\mathbf{C}$  of complex numbers and take their values in a topological vector space over  $\mathbf{C}$  (vector functions of a complex variable). Some of these definitions and properties extend even to functions which are defined on a subset of an arbitrary commutative *topological field*  $K$  and take their values in a topological vector space over  $K$ .

We shall indicate these generalizations in passing (see in particular I, p. 10, *Remark 2*), emphasising above all the case of functions of a complex variable, which are by far the most important, together with functions of a real variable, and will be studied in greater depth in a later Book.

#### 1. DERIVATIVE OF A VECTOR FUNCTION

**DEFINITION 1.** Let  $\mathbf{f}$  be a vector function defined on an interval  $I \subset \mathbf{R}$  which does not reduce to a single point. We say that  $\mathbf{f}$  is differentiable at a point  $x_0 \in I$  if

$$\lim_{x \rightarrow x_0, x \in I, x \neq x_0} \frac{\mathbf{f}(x) - \mathbf{f}(x_0)}{x - x_0}$$
 exists (in the vector space where  $\mathbf{f}$  takes its values); the value of this limit is called the *first derivative* (or simply the *derivative*) of  $\mathbf{f}$  at the point  $x_0$ , and it is denoted by  $\mathbf{f}'(x_0)$  or  $D\mathbf{f}(x_0)$ .

If  $\mathbf{f}$  is differentiable at the point  $x_0$ , so is the *restriction* of  $\mathbf{f}$  to any interval  $J \subset I$  which does not reduce to a single point and such that  $x_0 \in J$ ; and the derivative of this restriction is equal to  $\mathbf{f}'(x_0)$ . Conversely, let  $J$  be an interval contained in  $I$  and containing a neighbourhood of  $x_0$  relative to  $I$ ; if the restriction of  $\mathbf{f}$  to  $J$  admits a derivative at the point  $x_0$ , then so does  $\mathbf{f}$ .

We summarise these properties by saying that the concept of derivative is a *local* concept.

*Remarks.* \*1) In Kinematics, if the point  $\mathbf{f}(t)$  is the position of a moving point in the space  $\mathbf{R}^3$  at time  $t$ , then  $\frac{\mathbf{f}(t) - \mathbf{f}(t_0)}{t - t_0}$  is termed the *average velocity* between the instants  $t_0$  and  $t$ , and its limit  $\mathbf{f}'(t_0)$  is the *instantaneous velocity* (or simply *velocity*) at the time  $t_0$  (when this limit exists).\*

2) If a function  $\mathbf{f}$ , defined on  $I$ , is differentiable at a point  $x_0 \in I$ , it is necessarily *continuous relative to  $I$*  at this point.

**DEFINITION 2.** Let  $\mathbf{f}$  be a vector function defined on an interval  $I \subset \mathbf{R}$ , and let  $x_0$  be a point of  $I$  such that the interval  $I \cap [x_0, +\infty[$  (resp.  $I \cap ]-\infty, x_0]$ ) does not reduce to a single point. We say that  $\mathbf{f}$  is *differentiable on the right* (resp. *on the left*) at the point  $x_0$  if the restriction of  $\mathbf{f}$  to the interval  $I \cap [x_0, +\infty[$  (resp.  $I \cap ]-\infty, x_0]$ ) is differentiable at the point  $x_0$ ; the value of the derivative of this restriction at the point  $x_0$  is called the *right* (resp. *left*) derivative of  $\mathbf{f}$  at the point  $x_0$  and is denoted by  $\mathbf{f}'_d(x_0)$  (resp.  $\mathbf{f}'_g(x_0)$ ).

Let  $\mathbf{f}$  be a vector function defined on  $I$ , and  $x_0$  an *interior* point of  $I$  such that  $\mathbf{f}$  is continuous at this point; it follows from defs. 1 and 2 that for  $\mathbf{f}$  to be differentiable at  $x_0$  it is necessary and sufficient that  $\mathbf{f}$  admit both a right and a left derivative at this point, and that these derivatives be *equal*; and then

$$\mathbf{f}'(x_0) = \mathbf{f}'_d(x_0) = \mathbf{f}'_g(x_0).$$

*Examples.* 1) A *constant* function has zero derivative at every point.

2) An affine linear function  $x \mapsto \mathbf{a}x + \mathbf{b}$  has derivative equal to  $\mathbf{a}$  at every point.

3) The real function  $1/x$  (defined for  $x \neq 0$ ) is differentiable at each point  $x_0 \neq 0$ , for we have  $\left(\frac{1}{x} - \frac{1}{x_0}\right) / (x - x_0) = -\frac{1}{xx_0}$ , and, since  $1/x$  is continuous at  $x_0$ , the limit of the preceding expression is  $-1/x_0^2$ .

4) The scalar function  $|x|$ , defined on  $\mathbf{R}$ , has right derivative  $+1$  and left derivative  $-1$  at  $x = 0$ ; it is not differentiable at this point.

\*5) The real function equal to 0 for  $x = 0$ , and to  $x \sin 1/x$  for  $x \neq 0$ , is defined and continuous on  $\mathbf{R}$ , but has neither right nor left derivative at the point  $x \neq 0$ .\* One can give examples of functions which are continuous on an interval and fail to have a derivative at every point of the interval (I, p. 35, exerc. 2 and 3).

**DEFINITION 3.** We say that a vector function  $\mathbf{f}$  defined on an interval  $I \subset \mathbf{R}$  is *differentiable* (resp. *right differentiable*, *left differentiable*) on  $I$  if it is differentiable (resp. *right differentiable*, *left differentiable*) at each point of  $I$ ; the function  $x \mapsto \mathbf{f}'(x)$  (resp.  $x \mapsto \mathbf{f}'_d(x)$ ,  $x \mapsto \mathbf{f}'_g(x)$ ) defined on  $I$ , is called the *derived function*, or (by abuse of language) the *derivative* (resp. *right derivative*, *left derivative*) of  $\mathbf{f}$ , and is denoted by  $\mathbf{f}'$  or  $D\mathbf{f}$  or  $d\mathbf{f}/dx$  (resp.  $\mathbf{f}'_d$ ,  $\mathbf{f}'_g$ ).

*Remark.* A function may be differentiable on an interval without its derivative being continuous at every point of the interval (cf. I, p. 36, exerc. 5); \*this is shown by the

example of the function equal to 0 for  $x = 0$  and to  $x^2 \sin 1/x$  for  $x \neq 0$ ; it has a derivative everywhere, but this derivative is discontinuous at the point  $x = 0$ .\*

## 2. LINEARITY OF DIFFERENTIATION

**PROPOSITION 1.** *The set of vector functions defined on an interval  $I \subset \mathbf{R}$ , taking values in a given topological vector space  $E$ , and differentiable at the point  $x_0$ , is a vector space over  $\mathbf{R}$ , and the map  $\mathbf{f} \mapsto \mathbf{Df}(x_0)$  is a linear mapping of this space into  $E$ .*

In other words, if  $\mathbf{f}$  and  $\mathbf{g}$  are defined on  $I$  and differentiable at the point  $x_0$ , then  $\mathbf{f} + \mathbf{g}$  and  $\mathbf{f}a$  ( $a$  an arbitrary scalar) are differentiable at  $x_0$  and their derivatives there are  $\mathbf{f}'(x_0) + \mathbf{g}'(x_0)$  and  $\mathbf{f}'(x_0)a$  respectively. This follows immediately from the continuity of  $\mathbf{x} + \mathbf{y}$  and of  $\mathbf{x}a$  on  $E \times E$  and  $E$  respectively.

**COROLLARY.** *The set of vector functions defined on an interval  $I$ , taking values in a given topological vector space  $E$ , and differentiable on  $I$ , is a vector space over  $\mathbf{R}$ , and the map  $\mathbf{f} \mapsto \mathbf{Df}$  is a linear mapping of this space into the vector space of mappings from  $I$  into  $E$ .*

*Remark.* If one endows the vector space of mappings from  $I$  into  $E$  and its subspace of differentiable mappings (cf. *Gen. Top.*, X, p. 277) with the topology of simple convergence (or the topology of uniform convergence), the linear mapping  $\mathbf{f} \mapsto \mathbf{Df}$  is not continuous (in general) \*for example, the sequence of functions  $\mathbf{f}_n(x) = \sin n^2x/n$  converges uniformly to 0 on  $\mathbf{R}$ , but the sequence of derivatives  $\mathbf{f}'_n(x) = n \cos n^2x$  does not converge even simply to 0.\*

**PROPOSITION 2.** *Let  $E$  and  $F$  be two topological vector spaces over  $\mathbf{R}$ , and  $\mathbf{u}$  a continuous linear map from  $E$  into  $F$ . If  $\mathbf{f}$  is a vector function defined on an interval  $I \subset \mathbf{R}$ , taking values in  $E$ , and differentiable at the point  $x_0 \in I$ , then the composite function  $\mathbf{u} \circ \mathbf{f}$  has a derivative equal to  $\mathbf{u}(\mathbf{f}'(x_0))$  at  $x_0$ .*

Indeed, since 
$$\frac{\mathbf{u}(\mathbf{f}(x)) - \mathbf{u}(\mathbf{f}(x_0))}{x - x_0} = \mathbf{u} \left( \frac{\mathbf{f}(x) - \mathbf{f}(x_0)}{x - x_0} \right),$$
 this follows from the continuity of  $\mathbf{u}$ .

**COROLLARY.** *If  $\varphi$  is a continuous linear form on  $E$ , then the real function  $\varphi \circ \mathbf{f}$  has a derivative equal to  $\varphi(\mathbf{f}'(x_0))$  at the point  $x_0$ .*

*Examples.* 1) Let  $\mathbf{f} = (f_i)_{1 \leq i \leq n}$  be a function with values in  $\mathbf{R}^n$ , defined on an interval  $I \subset \mathbf{R}$ ; each real function  $f_i$  is none other than the composite function  $\text{pr}_i \circ \mathbf{f}$ , so is differentiable at the point  $x_0$  if  $\mathbf{f}$  is, and, if so,  $\mathbf{f}'(x_0) = (f'_i(x_0))_{1 \leq i \leq n}$ .

\*2) In Kinematics, if  $\mathbf{f}(t)$  is the position of a moving point  $M$  at time  $t$ , if  $\mathbf{g}(t)$  is the position at the same instant of the projection  $M'$  of  $M$  onto a plane  $P$  (resp. a line  $D$ ) with kernel a line (resp. a plane) not parallel to  $P$  (resp.  $D$ ), then  $\mathbf{g}$  is the composition of the projection  $\mathbf{u}$  of  $\mathbf{R}^3$  onto  $P$  (resp.  $D$ ) and of  $\mathbf{f}$ ; since  $\mathbf{u}$  is a (continuous) linear mapping

one sees that the projection of the velocity of a moving point onto a plane (resp. a line) is equal to the velocity of the projection of the moving point onto the plane (resp. line)\*.

3) Let  $f$  be a complex-valued function defined on an interval  $I \subset \mathbf{R}$ , and let  $a$  be an arbitrary complex number; prop. 2 shows that if  $f$  is differentiable at a point  $x_0$  then so is  $af$ , and the derivative of this function at  $x_0$  is equal to  $af'(x_0)$ .

### 3. DERIVATIVE OF A PRODUCT

Let us now consider  $p$  topological vector spaces  $E_i$  ( $1 \leq i \leq p$ ) over  $\mathbf{R}$ , and a continuous multilinear<sup>1</sup> map (which we shall denote by

$$(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_p) \mapsto [\mathbf{x}_1 \cdot \mathbf{x}_2 \dots \mathbf{x}_p]$$

of  $E_1 \times E_2 \times \dots \times E_p$  into a topological vector space  $F$  over  $\mathbf{R}$ .

**PROPOSITION 3.** For each index  $i$  ( $1 \leq i \leq p$ ) let  $\mathbf{f}_i$  be a function defined on an interval  $I \subset \mathbf{R}$ , taking values in  $E_i$ , and differentiable at the point  $x_0 \in I$ . Then the function

$$x \mapsto [\mathbf{f}_1(x) \cdot \mathbf{f}_2(x) \dots \mathbf{f}_p(x)]$$

defined on  $I$  with values in  $F$  has a derivative equal to

$$\sum_{i=1}^p [\mathbf{f}_1(x_0) \dots \mathbf{f}_{i-1}(x_0) \cdot \mathbf{f}'_i(x_0) \cdot \mathbf{f}_{i+1}(x_0) \dots \mathbf{f}_p(x_0)] \quad (1)$$

at  $x_0$ .

Let us put  $\mathbf{h}(x) = [\mathbf{f}_1(x) \cdot \mathbf{f}_2(x) \dots \mathbf{f}_p(x)]$ ; then, by the identity

$$[\mathbf{b}_1 \cdot \mathbf{b}_2 \dots \mathbf{b}_p] - [\mathbf{a}_1 \cdot \mathbf{a}_2 \dots \mathbf{a}_p] = \sum_{i=1}^p [\mathbf{b}_1 \dots \mathbf{b}_{i-1} \cdot (\mathbf{b}_i - \mathbf{a}_i) \cdot \mathbf{a}_{i+1} \dots \mathbf{a}_p],$$

we can write

$$\mathbf{h}(x) - \mathbf{h}(x_0) = \sum_{i=1}^p [\mathbf{f}_1(x) \dots \mathbf{f}_{i-1}(x) \cdot (\mathbf{f}_i(x) - \mathbf{f}_i(x_0)) \cdot \mathbf{f}_{i+1}(x_0) \dots \mathbf{f}_p(x_0)].$$

On multiplying both sides by  $\frac{1}{x - x_0}$  and letting  $x$  approach  $x_0$  in  $I$ , we obtain the expression (1), since both the map

$$(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_p) \mapsto [\mathbf{x}_1 \cdot \mathbf{x}_2 \dots \mathbf{x}_p]$$

and addition in  $F$  are continuous.

<sup>1</sup> Recall (*Alg.*, II, p. 265) that a map  $\mathbf{f}$  of  $E_1 \times E_2 \times \dots \times E_p$  into  $F$  is said to be *multilinear* if each partial mapping

$$\mathbf{x}_i \mapsto \mathbf{f}(\mathbf{a}_1, \dots, \mathbf{a}_{i-1}, \mathbf{x}_i, \mathbf{a}_{i+1}, \dots, \mathbf{a}_p)$$

from  $E_i$  into  $F$  ( $1 \leq i \leq p$ ) is a *linear* map, the  $\mathbf{a}_j$  for indices  $j \neq i$  being arbitrary in  $E_j$ . We note that if the  $E_i$  are *finite* dimensional over  $\mathbf{R}$  then every multilinear map of  $E_1 \times E_2 \times \dots \times E_p$  into  $F$  is necessarily *continuous*; this need not be so if some of these spaces are topological vector spaces of infinite dimension.

When some of the functions  $\mathbf{f}_i$  are *constant*, the terms in the expression (1) containing their derivatives  $\mathbf{f}'_i(x_0)$  are zero.

Let us consider in detail the particular case  $p = 2$ , the most important in applications: if  $(\mathbf{x}, \mathbf{y}) \mapsto [\mathbf{x}, \mathbf{y}]$  is a *continuous bilinear* map of  $E \times F$  into  $G$ , ( $E, F, G$  being topological vector spaces over  $\mathbf{R}$ ), and  $\mathbf{f}$  and  $\mathbf{g}$  are two vector functions, differentiable at  $x_0$ , with values in  $E$  and  $F$  respectively, then the vector function  $x \mapsto [\mathbf{f}(x), \mathbf{g}(x)]$  (which we denote by  $[\mathbf{f}, \mathbf{g}]$ ) has a derivative equal to  $[\mathbf{f}'(x_0), \mathbf{g}(x_0)] + [\mathbf{f}(x_0), \mathbf{g}'(x_0)]$  at  $x_0$ . In particular, if  $\mathbf{a}$  is a constant vector, then  $[\mathbf{a}, \mathbf{f}]$  (resp.  $[\mathbf{f}, \mathbf{a}]$ ) has a derivative equal to  $[\mathbf{a}, \mathbf{f}'(x_0)]$  (resp.  $[\mathbf{f}'(x_0), \mathbf{a}]$ ) at  $x_0$ .

If  $\mathbf{f}$  and  $\mathbf{g}$  are both differentiable on  $I$  then so is  $[\mathbf{f}, \mathbf{g}]$ , and we have

$$[\mathbf{f}, \mathbf{g}]' = [\mathbf{f}', \mathbf{g}] + [\mathbf{f}, \mathbf{g}'] \quad (2)$$

*Examples.* 1) Let  $f$  be a real function,  $\mathbf{g}$  a vector function, both differentiable at a point  $x_0$ ; the function  $\mathbf{g}f$  has a derivative equal to  $\mathbf{g}'(x_0)f(x_0) + \mathbf{g}(x_0)f'(x_0)$  at  $x_0$ . In particular, if  $\mathbf{a}$  is constant, then  $\mathbf{a}f$  has derivative  $\mathbf{a}f'(x_0)$ . This last remark, in conjunction with example 1 of I, p. 5, proves that if  $\mathbf{f} = (f_i)_{1 \leq i \leq n}$  is a vector function with values in  $\mathbf{R}^n$ , then for  $\mathbf{f}$  to be differentiable at the point  $x_0$  it is necessary and sufficient that each of the real functions  $f_i$  ( $1 \leq i \leq n$ ) be differentiable there: for, if  $(\mathbf{e}_i)_{1 \leq i \leq n}$  is the canonical basis of  $\mathbf{R}^n$ , we can write  $\mathbf{f} = \sum_{i=1}^n \mathbf{e}_i f_i$ .

2) The real function  $x^n$  arises from the multilinear function

$$(x_1, x_2, \dots, x_n) \mapsto x_1 x_2 \dots x_n$$

defined on  $\mathbf{R}^n$ , by substituting  $x$  for each of the  $x_i$ ; so prop. 3 shows that  $x^n$  is differentiable on  $\mathbf{R}$  and has derivative  $n x^{n-1}$ . As a result the polynomial function  $\mathbf{a}_0 x^n + \mathbf{a}_1 x^{n-1} + \dots + \mathbf{a}_{n-1} x + \mathbf{a}_n$  (the  $\mathbf{a}_i$  being constant vectors) has derivative

$$n \mathbf{a}_0 x^{n-1} + (n-1) \mathbf{a}_1 x^{n-2} + \dots + \mathbf{a}_{n-1};$$

when the  $\mathbf{a}_i$  are real numbers this function coincides with the derivative of a polynomial function as defined in Algebra (A, IV).

3) The euclidean *scalar product*  $(\mathbf{x} | \mathbf{y})$  (*Gen. Top.*, VI, p. 40) is a bilinear map (necessarily continuous) of  $\mathbf{R}^n \times \mathbf{R}^n$  into  $\mathbf{R}$ . If  $\mathbf{f}$  and  $\mathbf{g}$  are two vector functions with values in  $\mathbf{R}^n$ , and differentiable at the point  $x_0$ , then the real function  $x \mapsto (\mathbf{f}(x) | \mathbf{g}(x))$  has a derivative equal to  $(\mathbf{f}'(x_0) | \mathbf{g}(x_0)) + (\mathbf{f}(x_0) | \mathbf{g}'(x_0))$  at the point  $x_0$ . There is an analogous result for the hermitian scalar product on  $\mathbf{C}^n$ , this space being considered as a vector space over  $\mathbf{R}$ .

Let us consider in particular the case where the euclidean norm  $\|\mathbf{f}(x)\|$  is *constant*, so that  $(\mathbf{f}(x) | \mathbf{f}(x)) = \|\mathbf{f}(x)\|^2$  is also constant; on writing that the derivative of  $(\mathbf{f}(x) | \mathbf{f}(x))$  vanishes at  $x_0$  we obtain  $(\mathbf{f}'(x_0) | \mathbf{f}(x_0)) = 0$ ; in other words,  $\mathbf{f}'(x_0)$  is *orthogonal* to  $\mathbf{f}(x_0)$ .

4) If  $E$  is a *topological algebra* over  $\mathbf{R}$  (*cf.* Introduction), the product  $\mathbf{x}\mathbf{y}$  of two elements of  $E$  is a continuous bilinear function of  $(\mathbf{x}, \mathbf{y})$ ; if  $\mathbf{f}$  and  $\mathbf{g}$  have their values in  $E$  and are differentiable at the point  $x_0$ , then the function  $x \mapsto \mathbf{f}(x)\mathbf{g}(x)$  has a derivative equal to  $\mathbf{f}'(x_0)\mathbf{g}(x_0) + \mathbf{f}(x_0)\mathbf{g}'(x_0)$  at  $x_0$ . In particular, if  $U(x) = (\alpha_{ij}(x))$  and  $V(x) = (\beta_{ij}(x))$  are two *square matrices* of order  $n$ , differentiable at  $x_0$ , their product  $UV$  has a derivative equal to  $U'(x_0)V(x_0) + U(x_0)V'(x_0)$  at  $x_0$  (where  $U'(x) = (\alpha'_{ij}(x))$  and  $V'(x) = (\beta'_{ij}(x))$ ).

5) The *determinant*  $\det(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n)$  of  $n$  vectors  $\mathbf{x}_i = (x_{ij})_{1 \leq j \leq n}$  from the space  $\mathbf{R}^n$  (*Alg.*, III, p. 522) being a (continuous) multilinear function of the  $\mathbf{x}_i$ , one sees that if the

$n^2$  real functions  $f_{ij}$  are differentiable at  $x_0$ , then their determinant  $g(x) = \det(f_{ij}(x))$  has a derivative equal to

$$\sum_{i=1}^n \left[ \mathbf{f}_1(x_0), \dots, \mathbf{f}_{i-1}(x_0), \mathbf{f}'_i(x_0), \mathbf{f}_{i+1}(x_0), \dots, \mathbf{f}_n(x_0) \right]$$

at  $x_0$ , where  $\mathbf{f}_i(x) = (f_{ij}(x))_{1 \leq j \leq n}$ ; in other words, one obtains the derivative of a determinant of order  $n$  by taking the sum of the  $n$  determinants formed by replacing, for each  $i$ , the terms of the  $i^{\text{th}}$  column by their derivatives.

*Remark.* If  $U(x)$  is a square matrix which is differentiable and invertible at the point  $x_0$ , then the derivative of its determinant  $\Delta(x) = \det(U(x))$  can be expressed through the derivative of  $U(x)$  by the formula

$$\Delta'(x_0) = \Delta(x_0) \cdot \text{Tr}(U'(x_0)U^{-1}(x_0)). \quad (3)$$

Indeed, let us put  $U(x_0 + h) = U(x_0) + hV$ ; then, by definition,  $V$  tends to  $U'(x_0)$  when  $h$  tends to 0. One can write

$$\Delta(x_0 + h) = \Delta(x_0) \cdot \det(I + hVU^{-1}(x_0)).$$

Now  $\det(I + hX) = 1 + h\text{Tr}(X) + \sum_{k=2}^n \lambda_k h^k$ , the  $\lambda_k$  ( $k \geq 2$ ) being polynomials in the elements of the matrix  $X$ ; since the elements of  $VU^{-1}(x_0)$  have a limit when  $h$  tends to 0, we indeed obtain the formula (3).

#### 4. DERIVATIVE OF THE INVERSE OF A FUNCTION

**PROPOSITION 4.** *Let  $E$  be a complete normed algebra with a unit element over  $\mathbf{R}$  and let  $\mathbf{f}$  be a function defined on an interval  $I \subset \mathbf{R}$ , taking values in  $E$ , and differentiable at the point  $x_0 \in I$ . If  $\mathbf{y}_0 = \mathbf{f}(x_0)$  is invertible<sup>2</sup> in  $E$ , then the function  $x \mapsto (\mathbf{f}(x))^{-1}$  is defined on a neighbourhood of  $x_0$  (relative to  $I$ ), and has a derivative equal to  $-(\mathbf{f}(x_0))^{-1} \mathbf{f}'(x_0) (\mathbf{f}(x_0))^{-1}$  at  $x_0$ .*

Indeed, the set of invertible elements in  $E$  is an open set on which the function  $\mathbf{y} \mapsto \mathbf{y}^{-1}$  is continuous (*Gen. Top.*, IX, p. 178); since  $\mathbf{f}$  is continuous (relative to  $I$ ) at  $x_0$ ,  $(\mathbf{f}(x))^{-1}$  is defined on a neighbourhood of  $x_0$ , and we have

$$(\mathbf{f}(x))^{-1} - (\mathbf{f}(x_0))^{-1} = (\mathbf{f}(x))^{-1} (\mathbf{f}(x_0) - \mathbf{f}(x)) (\mathbf{f}(x_0))^{-1}.$$

The proposition thus follows from the continuity of  $\mathbf{y}^{-1}$  on a neighbourhood of  $\mathbf{y}_0$  and the continuity of  $\mathbf{xy}$  on  $E \times E$ .

<sup>2</sup> Recall from (*Alg.*, I, p. 15) that an element  $\mathbf{z} \in E$  is said to be *invertible* if there exists an element of  $E$ , denoted by  $\mathbf{z}^{-1}$ , such that  $\mathbf{zz}^{-1} = \mathbf{z}^{-1}\mathbf{z} = \mathbf{e}$  ( $\mathbf{e}$  being the unit element of  $E$ ).

*Examples.* 1) The most important particular case is that where  $E$  is one of the fields  $\mathbf{R}$  or  $\mathbf{C}$ : if  $f$  is a function with real or complex values, differentiable at the point  $x_0$ , and such that  $f(x_0) \neq 0$ , then  $1/f$  has derivative equal to  $-f'(x_0)/(f(x_0))^2$  at  $x_0$ .

2) If  $U = (\alpha_{ij}(x))$  is a square matrix of order  $n$ , differentiable at  $x_0$  and invertible at this point, then  $U^{-1}$  has derivative equal to  $-U^{-1}U'U^{-1}$  at  $x_0$ .

## 5. DERIVATIVE OF A COMPOSITE FUNCTION

**PROPOSITION 5.** *Let  $f$  be a real function defined on an interval  $I \subset \mathbf{R}$ , and  $\mathbf{g}$  a vector function defined on an interval of  $\mathbf{R}$  containing  $f(I)$ . If  $f$  is differentiable at the point  $x_0$  and  $\mathbf{g}$  is differentiable at the point  $f(x_0)$  then the composite function  $\mathbf{g} \circ f$  has a derivative equal to  $\mathbf{g}'(f(x_0))f'(x_0)$  at  $x_0$ .*

Let us put  $\mathbf{h} = \mathbf{g} \circ f$ ; for  $x \neq x_0$  we can write

$$\frac{\mathbf{h}(x) - \mathbf{h}(x_0)}{x - x_0} = \mathbf{u}(x) \frac{f(x) - f(x_0)}{x - x_0}$$

where we set  $\mathbf{u}(x) = \frac{\mathbf{g}(f(x)) - \mathbf{g}(f(x_0))}{f(x) - f(x_0)}$  if  $f(x) \neq f(x_0)$ , and  $\mathbf{u}(x) = \mathbf{g}'(f(x_0))$

otherwise. Now  $f(x)$  has limit  $f(x_0)$  when  $x$  tends to  $x_0$ , so  $\mathbf{u}(x)$  has limit  $\mathbf{g}'(f(x_0))$ , from which the proposition follows in view of the continuity of the function  $\mathbf{y}x$  on  $E \times \mathbf{R}$ .

## 6. DERIVATIVE OF AN INVERSE FUNCTION

**PROPOSITION 6.** *Let  $f$  be a homeomorphism of an interval  $I \subset \mathbf{R}$  onto an interval  $J = f(I) \subset \mathbf{R}$ , and let  $g$  be the inverse homeomorphism<sup>3</sup>. If  $f$  is differentiable at the point  $x_0 \in I$ , and if  $f'(x_0) \neq 0$ , then  $g$  has a derivative equal to  $1/f'(x_0)$  at  $y_0 = f(x_0)$ .*

For each  $y \in J$  we have  $g(y) \in I$  and  $u = f(g(y))$ ; we thus can write  $\frac{g(y) - g(y_0)}{y - y_0} = \frac{g(y) - x_0}{f(g(y)) - f(x_0)}$ , for  $y \neq y_0$ . When  $y$  tends to  $y_0$  while remaining in  $J$  and  $\neq y_0$ , then  $g(y)$  tends to  $x_0$  remaining in  $I$  and  $\neq x_0$ , and the right-hand side in the preceding formula thus has limit  $1/f'(x_0)$ , since by hypothesis  $f'(x_0) \neq 0$ .

**COROLLARY.** *If  $f$  is differentiable on  $I$  and if  $f'(x) \neq 0$  on  $I$ , then  $g$  is differentiable on  $J$  and its derivative at each point  $y \in J$  is  $1/f'(g(y))$ .*

For example, for each integer  $n > 0$ , the function  $x^{1/n}$  is a homeomorphism of  $\mathbf{R}_+$  onto itself, is the inverse of  $x^n$ , and has derivative  $\frac{1}{n}x^{\frac{1}{n}-1}$  at each  $x > 0$ .

One deduces easily, from prop. 5, that for every rational number  $r = p/q > 0$  the function  $x^r = (x^{1/q})^p$  has derivative  $rx^{r-1}$  at every  $x > 0$ .

<sup>3</sup> For  $f$  to be a homeomorphism of  $I$  onto a subset of  $\mathbf{R}$  we know that it is necessary and sufficient that  $f$  be continuous and strictly monotone on  $I$  (*Gen. Top.*, IV, p. 338, th. 5).

*Remarks.* 1) All the preceding propositions, stated for functions differentiable at a point  $x_0$ , immediately yield propositions for functions which are right (resp. left) differentiable at  $x_0$ , when, instead of the functions themselves, one considers their restrictions to the intersection of their intervals of definition with the interval  $[x_0, +\infty[$  (resp.  $] -\infty, x_0]$ ; we leave it to the reader to state them.

2) The preceding definitions and propositions (except for those concerning right and left derivatives) extend easily to the case where one replaces  $\mathbf{R}$  by an arbitrary *commutative non-discrete topological field*  $\mathbf{K}$ , and the topological vector spaces (resp. topological algebras) over  $\mathbf{R}$  by topological vector spaces (resp. topological algebras) over  $\mathbf{K}$ . In def. 1 and props. 1, 2 and 3 it is enough to replace  $I$  by a *neighbourhood* of  $x_0$  in  $\mathbf{K}$ ; in prop. 4 one must assume further that the map  $\mathbf{y} \mapsto \mathbf{y}^{-1}$  is defined and continuous on a neighbourhood of  $\mathbf{f}(x_0)$  in  $E$ . Prop. 5 generalizes in the following manner: let  $\mathbf{K}'$  be a non-discrete subfield of the topological field  $\mathbf{K}$ , let  $E$  be a topological vector space *over*  $\mathbf{K}$ ; let  $f$  be a function defined on a neighbourhood  $V \subset \mathbf{K}'$  of  $x_0 \in \mathbf{K}'$ , with values in  $\mathbf{K}$  (considered as a topological vector space over  $\mathbf{K}'$ ), differentiable at  $x_0$ , and let  $\mathbf{g}$  be a function defined on a neighbourhood of  $f(x_0) \in \mathbf{K}$ , with values in  $E$ , and differentiable at the point  $f(x_0)$ ; then the map  $\mathbf{g} \circ f$  is differentiable at  $x_0$  and has derivative  $\mathbf{g}'(f(x_0))f'(x_0)$  there ( $E$  being then considered as a topological vector space *over*  $\mathbf{K}'$ ).

With the same notation, let  $\mathbf{f}$  be a function defined on a neighbourhood  $V$  of  $a \in \mathbf{K}$ , with values in  $E$ , and differentiable at the point  $a$ ; if  $a \in \mathbf{K}'$ , then the *restriction* of  $\mathbf{f}$  to  $V \cap \mathbf{K}'$  is differentiable at  $a$ , and has derivative  $\mathbf{f}'(a)$  there. These considerations apply above all, in practice, to the case where  $\mathbf{K} = \mathbf{C}$  and  $\mathbf{K}' = \mathbf{R}$ .

Finally, prop. 6 extends to the case where one replaces  $I$  by a neighbourhood of  $x_0 \in \mathbf{K}$ , and  $f$  by a homeomorphism of  $I$  onto a neighbourhood  $J = f(I)$  of  $y_0 = f(x_0)$  in  $\mathbf{K}$ .

## 7. DERIVATIVES OF REAL-VALUED FUNCTIONS

The preceding definitions and propositions may be augmented in several respects when we deal with *real-valued* functions of a real variable.

In the first place, if  $f$  is such a function, defined on an interval  $I \subset \mathbf{R}$ , and continuous relative to  $I$  at a point  $x_0 \in I$ , it can happen that when  $x$  tends to  $x_0$  while remaining in  $I$  and  $\neq x_0$ , that  $\frac{f(x) - f(x_0)}{x - x_0}$  has a limit equal to  $+\infty$  or to  $-\infty$ ; one then says that  $f$  is differentiable at  $x_0$  and has derivative  $+\infty$  (resp.  $-\infty$ ) there; if the function  $f$  has a derivative  $f'(x)$  (finite or infinite) at every point  $x$  of  $I$ , then the function  $f'$  (with values in  $\overline{\mathbf{R}}$ ) is again called the *derived function* (or simply the *derivative*) of  $f$ . One generalizes the definitions of right and left derivative similarly.

*Example.* At the point  $x = 0$  the function  $x^{1/3}$  (the inverse function of  $x^3$ , a homeomorphism of  $\mathbf{R}$  onto itself) has a derivative, equal to  $+\infty$ ; at  $x = 0$  the function  $|x|^{1/3}$  has right derivative  $+\infty$  and left derivative  $-\infty$ .

The formulae for the derivative of a sum, of a product of differentiable real functions, and for the inverse of a differentiable function (props. 1, 3 and 4), as well as for the derivative of a (real-valued) composition of functions (prop. 5) remain valid when the derivatives that occur are infinite, so long as all the expressions that occur in these formulae make sense (*Gen. Top.*, IV, p. 345–346). In fact, if in prop. 6 one supposes that  $f$  is strictly increasing (resp. strictly decreasing) and continuous on  $I$ , and if  $f'(x_0) = 0$ , then the inverse function  $g$  has a derivative equal to  $+\infty$



- [\*\*read online Archaeology in Latin America pdf, azw \(kindle\)\*\*](#)
- [\*read online We Too Sing America: South Asian, Arab, Muslim, and Sikh Immigrants Shape Our Multiracial Future\*](#)
- [\*\*download online A Beautiful Question: Finding Nature's Deep Design\*\*](#)
- [read Archaeology in British Towns: From the Emperor Claudius to the Black Death](#)
- [Best Lesbian Romance 2013 pdf, azw \(kindle\)](#)
  
- <http://www.1973vision.com/?library/Chekhov.pdf>
- <http://www.celebritychat.in/?ebooks/We-Too-Sing-America--South-Asian--Arab--Muslim--and-Sikh-Immigrants-Shape-Our-Multiracial-Future.pdf>
- <http://bestarthritiscare.com/library/Too-Cool-for-This-School.pdf>
- <http://rodrigocaporal.com/library/Archaeology-in-British-Towns--From-the-Emperor-Claudius-to-the-Black-Death.pdf>
- <http://www.netc-bd.com/ebooks/The-Green-Pearl--Lyonesse--Book-2-.pdf>